Index of Authors

ACHARYA, A. S., MANJULA, B. N. & VITHAYATHIL, P. J. Structure and enzymic activity of ribonuclease-A esterified at glutamic acid-49 and aspartic acid-53 821–830

ADAMS, R. L. P. *see* BUTT, T. R. 309–314

ALHADEFF, J. A. Gel filtration of sialoglycoproteins 315–319

ALLEN, A. *see* MARSHALL, T. 569–578

ALLEN, G. & GREEN, N. M. Primary structures of cysteine-containing peptides from the calcium ion-transporting adenosine triphosphatase of rabbit sarcoplasmic reticulum 393–402

AL-SARRAJ, K., WHITE, D. A. & MAYER, R. J. Immunochemical characterization of casein from rabbit mammary gland 877–883

ANDREW, P. W. *see* TAKRURI, I. 459–466

ANTONINI, E. *see* GREENWOOD, C. 11–17

ARÁNYI, P. *see* MACHOVICH, R. 869–875

ASHCOM, J. *see* FRANZEN, J. S. 701–704

AXÉN, R. *see* CARLSSON, J. 723–737

BAINES, B. S. & BROCKLEHURST, K. A spectrophotometric method for the detection of contaminant chymopapains in preparations of papain. Selective modification of one type of thiol group in the chymopapains by a two-protonic-state reagent 345–347

BALDUINI, C. *see* SPEZIALE, P. 935–939

BANERJEE, C. K. *see* SANI, B. P. 643–649

BARBER, D., PARR, S. R. & GREENWOOD, C. The oxidation of *Pseudomonas* cytochrome *c*-551 oxidase by potassium ferricyanide 681–690

BARBER, D. *see also* GREENWOOD, C. 11–17

BARRETT, A. J. *see* WERB, Z. 949–958

BASHEY, H. M. *see* BASHEY, R. I. 885–894

BASHEY, R. I. BASHEY, H. M. & JIMENEZ, S. A. Characterization of pepsin-solubilized bovine heart-valve collagen 885–894

BASHEY, R. I. *see also* JIMENEZ, S. A. 337–340

BAXTER, A., CURRIE, L. M. & DURHAM, J. P. A general method for purification of deoxycytidine kinase 1005–1008

BEARPARK, T. M. & STIRLING, J. L. A difference in the specificities of human liver *N*-acetyl-β-hexosaminidases A and B detected by their activities towards glycosaminoglycan oligosaccharides 997–1000

BETTS, G. F. *see* BOSTIAN, K. A. 773–786, 787–798

BEYNON, R. J. & KAY, J. The inactivation of native enzyme by a neutral proteinase from rat intestinal muscle 291–298

BICKERSTAFF, G. F. & PRICE, N. C. Properties of matrix-bound dimer and monomer derivatives of immobilized creatine kinase from rabbit skeletal muscle 85–93

BOARD, P. G., MOORE, K. A. & SMITH, J. E. Purification and properties of γ-glutamylcyclotransferase from human erythrocytes 427–431

BOETTCHER, B. *see* KOLK, A. H. J. 767–771

BONNETT, R., CHARALAMBIDES, A. A., JONES, K., MAGNUS, I. A. & RIDGE, R. J. The direct determination of porphyrin carboxylic acids. High-pressure liquid chromatography with solvent systems containing phase-transfer agents 693–696

BOONYAPISIT, S. T. *see* OSTROW, J. D. 263–267

BOSTIAN, K. A. & BETTS, G. F. Kinetics and reaction mechanism of potassium-activated aldehyde dehydrogenase from *Saccharomyces cerevisiae* 787–798

BOSTIAN, K. A. & BETTS, G. F. Rapid purification and properties of potassium-activated aldehyde dehydrogenase from *Saccharomyces cerevisiae* 773–786

BOULTER, D. *see* TAKRURI, I. 459–466

BREWER, S. J. *see* TURNER, M. K. 839–850

BRITTAIN, T., GREENWOOD, C., SPRINGALL, J. P. & THOMSON, A. J. Magnetic-circular-dichroism studies of haem *a* and its derivatives 411–417

BROCKLEHURST, K. *see* BAINES, B. S. 345–347

BROWN, S. B. & DOCHERTY, J. C. Haem degradation in abnormal haemoglobins 985–987

BRUNORI, M. *see* GREENWOOD, C. 11–17

BRUSCHI, M. *see* MOURA, J. J. G. 419–425

BURCHELL, B. Substrate specificity and properties of uridine diphosphate glucuronyltransferase purified to apparent homogeneity from phenobarbital-treated rat liver 749–757

BURRIS, R. H. *see* LUDDEN, P. W. 1001–1003

BUTT, T. R., WOOD, W. M., MCKAY, E. L. & ADAMS, R. L. P. Involvement of deoxyribonucleic acid polymerase β in nuclear deoxyribonucleic acid synthesis 309–314

CAMMACK, R. *see* MOURA, J. J. G. 419–425

CARLSSON, J., DREVIN, H. & AXÉN, R. Protein thiolation and reversible protein–protein conjugation. *N*-Succinimidyl 3-(2-pyridyldithio)propionate, a new heterobifunctional reagent 723–737

CARNE, A. & MOORE, C. H. The amino acid sequence of the tryptic peptides from actinidin, a proteolytic enzyme from the fruit of *Actinidia chinensis* 73–83

CARRELL, R. W. *see* FRENCH, J. K. 19–26

CARROLL, M. Characterization of proteins structurally related to human *N*-acetyl-β-D-glucosaminidase 191–196

CHANCE, E. M. *see* CLORE, G. M. 799–810, 811–820

CHARALAMBIDES, A. A. *see* BONNETT, R. 693–696

CHELL, R. M., SUNDARAM, T. K. & WILKINSON, A. E. Isolation and characterization of isocitrate lyase from a thermophilic *Bacillus* sp. 165–177

CHIEN, J.-L. *see* LI, S.-C. 697–699

CHILDS, R. A., FEIZI, T., FUKUDA, M. & HAKOMORI, S.-I. Blood-group-I activity associated with Band 3, the major intrinsic membrane protein of human erythrocytes 333–336

CHILDS, R. A. *see also* FEIZI, T. 245–254

CHOONG, Y. S., SHEPHERD, M. G. & SULLIVAN, P. A. Inactivation and modification of lactate oxidase with fluorodinitrobenzene 255–262

CHRISTELLER, J. T. & LAING, W. A. A kinetic study of ribulose bisphosphate carboxylase from the photosynthetic bacterium *Rhodospirillum rubrum* 467–473

CLARKE, N. & DAWSON, R. M. C. *rac*-Glycerol 1:2-cyclic phosphate 2-phosphodiesterase, a new soluble phosphodiesterase of mammalian tissues 579–589

CLORE, G. M. & CHANCE, E. M. The mechanism of reaction of ferricyanide-pretreated mixed-valence-state membrane-bound cytochrome oxidase with oxygen at 173K 811–820

CLORE, G. M. & CHANCE, E. M. The mechanism of reaction of fully reduced membrane-bound cytochrome oxidase with oxygen at 176K 799–810
COLOSIMO, A. *see* GREENWOOD, C. 11–17
COOPER, C. G. *see* NARAYANAN, A. S. 857–862
CRAIG, R. K., MCILREAVY, D. & HALL, R. L. Separation and partial characterization of guinea-pig caseins 633–641
CRANG, A. J. & RUMSBY, M. G. Solubilization of isolated central-nervous-system myelin preparations by the anionic detergent sodium dodecyl sulphate 909–917
CREWTHER, W. G., INGLIS, A. S. & MCKERN, N. M. Amino acid sequences of α-helical segments from *S*-carboxymethylkerateine-A. Complete sequence of a type-II segment 365–371
CREWTHER, W. G. *see also* ELLEMAN, T. C. 387–391; GOUGH, K. H. 373–385; HOGG, D. MCC. 353–363
CURRIE, L. M. *see* BAXTER, A. 1005–1008

DARVEY, I. G. *see* WALKER, E. J. 1–4, 5–10
DAVIS, A. W. & PHILLIPS, D. R. A defined molecular-weight distribution of deoxyribonucleic acid after extensive sonication 179–183
DAWSON, R. M. C. *see* CLARKE, N. 579–589
DEAN, P. D. G. *see* WATSON, D. H. 591–596
DINGLE, J. T. *see* WERB, Z. 949–958
DIXON, H. B. F. *see* SARKAR, B. 895–897
DOCHERTY, J. C. *see* BROWN, S. B. 985–987
DOWLING, L. M. *see* HOGG, D.MCC. 353–363
DREVIN, H. *see* CARLSSON, J. 723–737
DUNN, M. J., KEMP, R. B. & MADDY, A. H. The similarity of the two high-molecular-weight polypeptides of erythrocyte spectrin 197–205
DURHAM, J. P. *see* BAXTER, A. 1005–1008
DWEK, R. A. *see* GETTINS, P. 713–722

EADY, R. R. *see* LOWE, D. J. 277–290
EDWARDS, M. R. *see* SHARP, J. A. 759–765
ELLEMAN, T. C., CREWTHER, W. G. & VAN DER TOUW, J. Amino acid sequences of α-helical segments from *S*-carboxymethylkerateine-A. Statistical analysis 387–391
ENGASSER, J.-M. & HISLAND, P. Diffusional increase and decrease in half-maximal-activity substrate concentrations with two-substrate enzymic reactions 341–343
EVANS, R. W. & WILLIAMS, J. Studies of the binding of different iron donors to human serum transferrin and isolation of iron-binding fragments from the *N*- and *C*-terminal regions of the protein 543–552
EVANS, R. W. *see also* WILLIAMS, J. 533–539

FARTHING, J. E. *see* TURNER, M. K. 839–850
FEINGOLD, D. S. *see* FRANZEN, J. S. 701–704
FEIZI, T., CHILDS, R. A., HAKOMORI, S.-I. & POWELL, M. E. Blood-group-Ii-active gangliosides of human erythrocyte membranes 245–254
FEIZI, T. *see also* CHILDS, R. A. 333–336
FISCHER, J., KOHLER, T. R., LIPSON, L. G., FLORES, J., WITKUM, P. A. & SHARP, G. W. G. Studies on the time course and rate-limiting steps in the activation of adenylate cyclase in rat liver by cholera toxin 59–64
FLANAGAN, P. R. & FORSTNER, G. G. Purification of rat intestinal maltase/glucoamylase and its anomalous dissociation either by heat or by low pH 553–563
FLORES, J. *see* FISCHER, J. 59–64
FORSTNER, G. G. *see* FLANAGAN, P. R. 553–563
FOSTER, J. A. *see* MECHAM, R. P. 617–625
FOWERAKER, A. R. *see* ISLES, M. 237–243
FRANZEN, J. S., MARCHETTI, P., ISHMAN, R., ASHCOM, J. & FEINGOLD, D. S. Half-sites oxidation of bovine liver uridine diphosphate glucose dehydrogenase 701–704
FRENCH, J. K., WINTERBOURN, C. C. & CARRELL, R. W. Mechanism of oxyhaemoglobin breakdown on reaction with acetylphenylhydrazine 19–26
FUKUDA, M. *see* CHILDS, R. A. 333–336

GALLIGANI, L. *see* SPEZIALE, P. 935–939
GARRICK, L. M., SLOAN, R. L., RYAN, T. W., KLONOWSKI, T. J. & GARRICK, M. D. Primary structure of the major β-chain of rat haemoglobins 321–330
GARRICK, M. D. *see* GARRICK, L. M. 321–330
GETTINS, P., GIVOL, D. & DWEK, R. A. The role of nitro groups in the binding of nitroaromatics to protein MOPC 315 713–722
GILON, C. *see* MUSZKAT, K. A. 993–996
GIVOL, D. *see* GETTINS, P. 713–722
GODDARD, J. P. *see* KELLY, J. M. 521–532
GOODWIN, G. H. *see* RABBANI, A. 497–505
GOUGH, K. H., INGLIS, A. S. & CREWTHER, W. G. Amino acid sequences of α-helical segments from *S*-carboxymethylkerateine-A. Complete sequence of a type-I segment 373–385
GRÄSBECK, R. *see* MARCOULLIS, G. 705–712
GREEN, N. M. *see* ALLEN, G. 393–402
GREENWOOD, C., BARBER, D., PARR, S. R., ANTONINI, E., BRUNORI, M. & COLOSIMO, A. The reaction of *Pseudomonas aeruginosa* cytochrome *c*-551 oxidase with oxygen 11–17
GREENWOOD, C. *see also* BARBER, D. 681–690; BRITTAIN, T. 411–417
GRUNNET, I. & KNUDSEN, J. Molecular weight and subunit size of rabbit mammary-gland fatty acid synthetase 929–933

HAKOMORI, S.-I. *see* CHILDS, R. A. 333–336; FEIZI, T. 245–254
HALL, D. O. *see* MOURA, J. J. G. 419–425
HALL, P. K. & ROBERTS, R. C. Physical and chemical properties of human plasma α_2-macroglobulin 27–38
HALL, R. L. *see* CRAIG, R. K. 633–641
HARDINGHAM, T. *see* ISLES, M. 237–243
HARDMAN, M. J. *see* PARKER, D. M. 269–275
HARVEY, M. J. *see* WATSON, D. H. 591–596
HASLETT, B. G. *see* TAKRURI, I. 459–466
HAYZER, D. J. & MOSES, V. Proline biosynthesis by cell-free extracts of *Escherichia coli* and potential errors arising from the use of a bioradiological assay procedure 207–217
HAYZER, D. J. & MOSES, V. The enzymes of proline biosynthesis in *Escherichia coli*. Their molecular weights and the problem of enzyme aggregation 219–228
HEMMILÄ, I. A. MÄNTSÄLÄ, P. I. Inactivation of glutamate dehydrogenase and glutamate synthesis from *Bacillus megaterium* by phenylglyoxal, butane-2,3-dione and pyridoxal 5′-phosphate 53–58

HEMMILÄ, I. A. & MÄNTSÄLÄ, P. I. Purification and properties of glutamate synthase and glutamate dehydrogenase from *Bacillus megaterium* 45–52
HILDEBRANDT, V. *see* NICHOLLS, P. 65–72
HISLAND, P. *see* ENGASSER, J.-M. 341–343
HOGG, D.McC., DOWLING, L. M. & CREWTHER, W. G. Amino acid sequences of α-helical segments from *S*-carboxymethylkerateine-A. Tryptic and chymotryptic peptides from a type-II-segment 353–363
HOLBROOK, J. J. *see* LODOLA, A. 597–605; PARKER, D. M. 269–275, 959–967
HRADEC, J. & KŘÍŽ, O. Heparin–Sepharose 4B at low temperatures retains ribosomes 349–352
HUANG, C. *see* MATSUO, M. 441–447
HUANG, L. C. *see* MATSUO, M. 441–447
HUI, G. *see* PEGG, A. E. 739–748
HUTSON, N. J. KERBEY, A. L., RANDLE, P. J. & SUGDEN, P. H. Conversion of inactive (phosphorylated) pyruvate dehydrogenase complex into active complex by the phosphate reaction in heart mitochondria is inhibited by alloxan-diabetes or starvation in the rat 669–680

INGLIS, A. S. *see* CREWTHER, W. G. 365–371; GOUGH, K. H. 373–385
ISHMAN, R. *see* FRANZEN, J. S. 701–704
ISLES, M., FOWERAKER, A. R., JENNINGS, B. R., HARDINGHAM, T. & MUIR, H. Characterization of proteoglycan and the proteoglycan–hyaluronic acid complex by electric birefringence 237–243

JACOBS, A. *see* WAGSTAFF, M. 969–977
JECK, R. *see* LODOLA, A. 597–605
JENNINGS, B. R. *see* ISLES, M. 237–243
JIMENEZ, S. A. & BASHEY, R. I. Solubilization of bovine heart-valve collagen 337–340
JIMENEZ, S. A. *see also* BASHEY, R. I. 885–894
JOHNS, E. W. *see* RABBANI, A. 497–505
JOHNSON, L. N. *see* PERKINS, S. J. 607–616
JONES, K. *see* BONNETT, R. 693–696
JONES, M. *see* THOMAS, P. 981–983
JONES, R. W. Histone composition of a chromatin fraction containing ribosomal deoxyribonucleic acid isolated from the macronucleus of *Tetrahymena pyriformis* 155–164
JONES, R. W. Preparation of chromatin containing ribosomal deoxyribonucleic acid from the macronucleus of *Tetrahymena pyriformis* 145–153

KAY, J. *see* BEYNON, R. J. 291–298
KELLY, J. M., GODDARD, J. P. & MADEN, B. E. H. Evidence on the conformation of HeLa-cell 5.8S ribosomal ribonucleic acid from the reaction of specific cytidine residues with sodium bisulphite 521–532
KEMP, R. B. *see* DUNN, M. J. 197–205
KERBEY, A. L. *see* HUTSON, N. J. 669–680
KLONOWSKI, T. J. *see* GARRICK, L. M. 321–330
KNUDSEN, J. *see* GRUNNET, I. 929–933
KOHLER, T. R. *see* FISCHER, J. 59–64
KOLK, A. H. J., VAN KUYK, L. & BOETTCHER, B. Isolation of human lactate dehydrogenase isoenzyme X by affinity chromatography 767–771
KŘÍŽ, O. *see* HRADEC, J. 349–352
KUZAN, F. *see* NARAYANAN, A. S. 857–862

LAING, W. A. *see* CHRISTELLER, J. T. 467–473
LAWSON, D. E. M. *see* WILSON, P. W. 627–631
LE GALL, J. *see* MOURA, J. J. G. 419–425
LEE, J. S. & WARING, M. J. Bifunctional intercalation and sequence specificity in the binding of quinomycin and triostin antibiotics to deoxyribonucleic acid 115–128
LEE, J. S. & WARING, M. J. Interaction between synthetic analogues of quinoxaline antibiotics and nucleic acids. Changes in mechanism and specificity related to structural alterations 129–144
LI, S.-C., CHIEN, J.-L., WAN, C. C. & LI, Y.-T. Occurrence of glycosphingolipids in chicken egg yolk 697–699
LI, Y.-T. *see* LI, S.-C. 697–699
LIPSON, L. G. *see* FISCHER, J. 59–64
LODOLA, A., PARKER, D. M., JECK, R. & HOLBROOK, J. J. Malate dehydrogenase of the cytosol. Ionizations of the enzyme–reduced-coenzyme complex and a comparison with lactate dehydrogenase 597–605
LODOLA, A. *see also* PARKER, D. M. 959–967
LOWE, D. J., EADY, R. R. & THORNELEY, R. N. F. Electron-paramagnetic-resonance studies on nitrogenase of *Klebsiella pneumoniae*. Evidence for acetylene- and ethylene–nitrogenase transient complexes 277–290
LUDDEN, P. W., OKON, Y. & BURRIS, R. H. The nitrogenase system of *Spirillum lipoferum* 1001–1003
LYNN, W. S. *see* SAHU, S. 565–568

MACHIN, P. A. *see* PERKINS, S. J. 607–616
MACHOVICH, R. & ARÁNYI, P. Effect of heparin on thrombin inactivation by antithrombin-III 869–875
MADDY, A. H. *see* DUNN, M. J. 197–205
MADEN, B. E. H. *see* KELLY, J. M. 521–532
MAGNUS, I. A. *see* BONNETT, R. 693–696
MANJULA, B. N. *see* ACHARYA, A. S. 821–830
MÄNTSÄLÄ, P. I. *see* HEMMILÄ, I. A. 45–52, 53–58
MARCHETTI, P. *see* FRANZEN, J. S. 701–704
MARCOULLIS, G., MERIVUORI, H. & GRÄSBECK, R. Comparative studies on intrinsic factor and cobalophilin in different parts of the gastrointestinal tract of the pig 705–712
MARSHALL, T. & ALLEN, A. The isolation and characterization of the high-molecular weight-glycoprotein from pig colonic mucus 569–578
MATSUO, M., HUANG, C. & HUANG, L. C. Evidence for an essential arginine recognition site on adenosine 3′:5′-cyclic monophosphate-dependent protein kinase of rabbit skeletal muscle 441–447
MATTHIEU, J.-M. Proteins from sciatic-nerve myelin in Quaking and Jimpy mice 989–991
MAYER, R. J. *see* AL-SARRAJ, K. 877–883
MCILREAVY, D. *see* CRAIG, R. K. 633–641
MCKAY, E. L. *see* BUTT, T. R. 309–314
MCKERN, N. M. *see* CREWTHER, W. G. 365–371
MECHAM, R. P. & FOSTER, J. A. A structural model for desmosine cross-linked peptides 617–625
MERIVUORI, H. *see* MARCOULLIS, G. 705–712
MERRETT, M. J. *see* WOODCOCK, E. 95–101
MILNE, J. G. C. Derivation of equations relating the fractional saturation of *N*-terminal and *C*-terminal sites to the number of bound iron atoms per molecule of ovotransferrin 541
MOORE, C. H. *see* CARNE, A. 73–83

MOORE, G. Influence of the peptide-chain length on disulphide-bond formation in neurohypophysial hormones and analogues 403–409
MOORE, K. A. *see* BOARD, P. G. 427–431
MORETON, K. *see* WILLIAMS, J. 533–539
MOSES, V. *see* HAYZER, D. J. 207–217, 219–228
MOURA, J. J. G., XAVIER, A. V., CAMMACK, R., HALL D. O., BRUSCHI, M. & LE GALL, J. Oxidation–reduction studies of the Mo–(2Fe–2S) protein from *Desulfovibrio gigas* 419–425
MUIR, H. *see* ISLES, M. 237–243
MUKKUR, T. K. S. Thermodynamics for the interaction of ε-dinitrophenyl-L-lysine and bovine colostral anti-dinitrophenyl immunoglobulin G2 39–44
MURRAY, R. K. *see* NARASIMHAN, R. 475–482
MUSZKAT, K. A., WEINSTEIN, M. & GILON, C. Biochemical applications of chemically induced nuclear polarization in phenols, peptides, catecholamines and related molecules 993–996

NARASIMHAN, R. & MURRAY, R. K. Comparative study of the glycosphingolipids of chicken bursa of Fabricius and of chicken, rat and human thymus 475–482
NARAYANAN, A. S., PAGE, R. C., KUZAN, F. & COOPER, C. G. Elastin cross-linking *in vitro*. Studies on factors influencing the formation of desmosines by lysyl oxidase action on tropoelastin 857–862
NICHOL, L. W. *see* TELLAM, R. 185–190
NICHOLLS, P. & HILDEBRANDT, V. Binding of ligands and spectral shifts in cytochrome *c* oxidase 65–72
NISHIKURA, K. Identification of histidine-122α in human haemoglobin as one of the unknown alkaline Bohr groups by hydrogen–tritium exchange 651–657

O'DONNELL, M. J. & SMITH, B. E. Electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein of nitrogenase between +50 and −450mV 831–838
OKON, Y. *see* LUDDEN, P. W. 1001–1003
OSTROW, J. D. & BOONYAPISIT, S. T. Inaccuracies in measurement of conjugated and unconjugated bilirubin in bile with ethyl anthranilate diazo and solvent-partition methods 263–267

PAGE, M. A. & TUBBS, P. K. Some properties of 3-hydroxy-3-methylglutaryl-coenzyme A synthase from ox liver 925–928
PAGE, R. C. *see* NARAYANAN, A. S. 857–862
PARKER, D. M., HARDMAN, M. J., PLAPP, B. V., HOLBROOK, J. J. & SHORE, J. D. pH-dependent changes of intrinsic fluorescence of chemically modified liver alcohol dehydrogenases 269–275
PARKER, D. M., LODOLA, A. & HOLBROOK, J. J. Use of the sulphite adduct of nicotinamide–adenine dinucleotide to study ionizations and the kinetics of lactate dehydrogenase and malate dehydrogenase 959–967
PARKER, D. M. *see also* LODOLA, A. 597–605
PARLAMENTI, R. *see* PETRUCCI, T. 229–235
PARR, S. R. *see* BARBER, D. 681–690; GREENWOOD, C. 11–17
PEGG, A. E. & HUI, G. Formation and subsequent removal of O^6-methylguanine from deoxyribonucleic acid in rat liver and kidney after small doses of dimethylnitrosamine 739–748
PENNINGS, E. J. M., VRIELINK, R. & VAN KEMPEN, G. M. J. Kinetics and mechanism of the rat brain phenol sulphotransferase reaction 299–307
PERKINS, S. J., JOHNSON, L. N., MACHIN, P. A. & PHILLIPS, D. C. Crystal structures of egg-white lysozyme of hen in acetate-free medium and of lysozyme complexes with *N*-acetylglucosamine and β-methyl *N*-acetylglucosaminide 607–616
PERRY, S. V. *see* WEEKS, R. A. 449–457
PETRUCCI, T., SANNIA, G., PARLAMENTI, R. & SILANO, V. Structural studies of wheat monomeric and dimeric protein inhibitors of α-amylase 229–235
PHILLIPS, D. C. *see* PERKINS, S. J. 607–616
PHILLIPS, D. R. *see* DAVIS, A. W. 179–183
PLAPP, B. V. *see* PARKER, D. M. 269–275
PLUMMER, D. T. *see* REAVILL, C. A. 851–856
POWELL, M. E. *see* FEIZI, T. 245–254
PRICE, N. C. *see* BICKERSTAFF, G. F. 85–93

RABBANI, A., GOODWIN, G. H. & JOHNS, E. W. Studies on the tissue specificity of the high-mobility-group non-histone chromosomal proteins from calf 497–505
RALSTON, G. B. *see* WALKER, E. J. 1–4, 5–10
RANDLE, P. J. *see* HUTSON, N. J. 669–680; SUGDEN, P. H. 659–668
REAVILL, C. A., WOOSTER, M. S. & PLUMMER, D. T. The interaction of purified acetylcholinesterase from pig brain with liposomes 851–856
REID, K. B. M. & THOMPSON, E. O. P. Amino acid sequence of the *N*-terminal 108 amino acid residues of the B chain of subcomponent C1q of the first component of human complement 863–868
REYNOLDS, J. J. *see* WERB, Z. 949–958
RIDGE, R. J. *see* BONNETT, R. 693–696
ROBERTS, G. P. Structural studies on the glycoproteins from bovine cervical mucus 941–947
ROBERTS, R. C. *see* HALL, P. K. 27–38
ROGERS, L. J. *see* TAKRURI, I. 459–466
ROGIE, A. *see* SKINNER, E. R. 507–520
RUMSBY, M. G. *see* CRANG, A. J. 909–917
RYAN, T. W. *see* GARRICK, L. M. 321–330

SAHU, S. & LYNN, W. S. Hyaluronic acid in the pulmonary secretions of patients with asthma 565–568
SANI, B. P. & BANERJEE, C. K. Purification and properties of retinoic acid-binding protein from chick-embryo skin 643–649
SANNIA, G. *see* PETRUCCI, T. 229–235
SARKAR, B., DIXON, H. B. F. & WEBSTER, D. Removal by transamination and scission of residues from the peptide representing the copper-transport site of serum albumin 895–897
SCOTT, J. E. & TIGWELL, M. J. Periodate oxidation and the shapes of glycosaminoglycuronans in solution 103–114
SHARP, G. W. G. *see* FISCHER, J. 59–64
SHARP, J. A. & EDWARDS, M. R. Purification and properties of succinyl-coenzyme A–3-oxo acid coenzyme A-transferase from sheep kidney 759–765
SHEPHERD, M. G. *see* CHOONG, Y. S. 255–262
SHORE, J. D. *see* PARKER, D. M. 269–275
SILANO, V. *see* PETRUCCI, T. 229–235

SKINNER, E. R. & ROGIE, A. The isolation and partial characterization of the serum lipoproteins and apolipoproteins of the rainbow trout 507–520
SLOAN, R. L. *see* GARRICK, L. M. 321–330
SMITH, B. E. *see* O'DONNELL, M. J. 831–838
SMITH, J. E. *see* BOARD, P. G. 427–431
SPEZIALE, M. S. *see* SPEZIALE, P. 935–939
SPEZIALE, P., SPEZIALE, M. S., GALLIGANI, L. & BALDUINI, C. Interactions between different corneal proteoglycans 935–939
SPRINGALL, J. P. *see* BRITTAIN, T. 411–417
STANSFIELD, D. A. *see* YOUNG, J. L. 919–924
STIRLING, J. L. *see* BEARPARK, T. M. 997–1000
SUGDEN, P. H. & RANDLE, P. J. Regulation of pig heart pyruvate dehydrogenase by phosphorylation. Studies on the subunit and phosphorylation stoicheiometries 659–668
SUGDEN, P. H. *see also* HUTSON, N. J. 669–680
SULLIVAN, P. A. *see* CHOONG, Y. S. 255–262
SUNDARAM, T. K. *see* CHELL, R. M. 165–177

TAKRURI, I., HASLETT, B. G., BOULTER, D., ANDREW, P. W. & ROGERS, L. J. The amino acid sequence of ferredoxin from the red alga *Porphyra umbilicalis* 459–466
TELLAM, R., WINZOR, D. J. & NICHOL, L. W. The role of zinc in the stabilization of the dimeric form of bacterial α-amylase 185–190
THOMAS, P. & JONES, M. The effects of Triton WR1339 and asialo-fetuin on the hepatic uptake of circulating native and asialo-carcinoembryonic antigen 981–983
THOMPSON, E. O. P. *see* REID, K. B. M. 863–868
THOMPSON, A. J. *see* BRITTAIN, T. 411–417
THORNELEY, R. N. F. *see* LOWE, D. J. 277–290
TIGWELL, M. J. *see* SCOTT, J. E. 103–114
TSAI, C. S. Kinetic and mechanistic studies of methylated liver alcohol dehydrogenase 483–496
TUBBS, P. K. *see* PAGE, M. A. 925–928
TURNER, M. K., FARTHING, J. E. & BREWER, S. J. The oxygenation of [3-*methyl*-^{3}H]desacetoxycephalosporin C [7β-(5-D-aminoadipamido)-3-methylceph-3-em-4-carboxylic acid] to [3-*hydroxymethyl*-^{3}H]desacetylcephalosporin C by 2-oxoglutarate-linked dioxygenases from *Acremonium chrysogenum* and *Streptomyces clavuligerus* 839–850

VAN DER TOUW, J. *see* ELLEMAN, T. C. 387–391
VAN KEMPEN, G. M. J. *see* PENNINGS, E. J. M. 299–307
VAN KUYK, L. *see* KOLK, A. H. J. 767–771
VITHAYATHIL, P. J. *see* ACHARYA, A. S. 821–830
VRIELINK, R. *see* PENNINGS, E. J. M. 299–307

WAGSTAFF, M., WORWOOD, M. & JACOBS, A. Properties of human tissue isoferritins 969–977
WALKER, E. J., RALSTON, G. B. & DARVEY, I. G. Preparation of allosteric ribonuclease 5–10
WALKER, E. J., RALSTON, G. B. & DARVEY, I. G. The nature of the allosteric interactions of ribonuclease and its ligands 1–4
WAN, C. C. *see* LI, S.-C. 697–699
WARING, M. J. *see* LEE, J. S. 115–128, 129–144
WATSON, D. H., HARVEY, M. J. & DEAN, P. D. G. The selective retardation of $NADP^+$-dependent dehydrogenases by immobilized Procion Red HE-3B 591–596
WATTS, C. & WHEELER, K. P. Protein and lipid components of the pigeon erythrocyte membrane 899–907
WEBSTER, D. *see* SARKAR, B. 895–897
WEEKS, R. A. & PERRY, S. V. Characterization of a region of the primary sequence of troponin C involved in calcium ion-dependent interaction with troponin I 449–457
WEINSTEIN, M. *see* MUSZKAT, K. A. 993–996
WERB, Z., DINGLE, J. T., REYNOLDS, J. J. & BARRETT, A. J. Proteoglycan-degrading enzymes of rabbit fibroblasts and granulocytes 949–958
WHEELER, K. P. *see* WATTS, C. 899–907
WHITE, D. A. *see* AL-SARRAJ, K. 877–883
WILKINSON, A. E. *see* CHELL, R. M. 165–177
WILLCOX, P. Secretion of β-*N*-acetylglucosaminidase isoenzymes by normal human fibroblasts 433–439
WILLIAMS, J., EVANS, R. W. & MORETON, K. The iron-binding properties of hen ovotransferrin 533–539
WILLIAMS, J. *see also* EVANS, R. W. 543–552
WILSON, P. W. & LAWSON, D. E. M. Incorporation of [^{3}H]leucine into an actin-like protein in response to 1,25-dihydroxycholecalciferol in chick intestinal brush borders 627–631
WINTERBOURN, C. C. *see* FRENCH, J. K. 19–26
WINZOR, D. J. *see* TELLAM, R. 185–190
WITKUM, P. A. *see* FISCHER, J. 59–64
WOOD, W. M. *see* BUTT, T. R. 309–314
WOODCOCK, E. & MERRETT, M. J. Purification and immunochemical characterization of malate synthase from *Euglena gracilis* 95–101
WOOSTER, M. S. *see* REAVILL, C. A. 851–856
WORWOOD, M. *see* WAGSTAFF, M. 969–977

XAVIER, A. V. *see* MOURA, J. J. G. 419–425

YOUNG, J. L. & STANSFIELD, D. A. Solubilization of bovine corpus-luteum adenylate cyclase in Lubrol-PX, Triton X-100 or digitonin and the stabilizing effect of sodium fluoride present in the solubilization medium 919–924

Index of Subjects

Acetylcholinesterase, brain, pig, purified, interaction of, with liposomes (Reavill, C. A., Wooster, M. S. & Plummer, D. T.) 851–856

Acetylene, electron-paramagnetic-resonance evidence for the formation of transient complexes of ethylene and, with *Klebsiella pneumoniae* nitrogenase complex (Lowe, D. J., Eady, R. R. & Thorneley, R. N. F.) 277–290

N-Acetylglucosamine, structures of crystals of chicken egg-white lysozyme and of complexes of the enzyme with β-methyl *N*-acetylglucosaminide and, grown in acetate-free medium (Perkins, S. J., Johnson, L. N., Machin, P. A. & Phillips, D. C.) 607–616

N-Acetyl-β-D-glucosaminidase, placenta, human, immunochemical characterization of human liver proteins structurally related to (Carroll, M.) 191–196

β-*N*-Acetylglucosaminidase, secretion of isoenzymes of, by cultured human skin fibroblasts (Willcox, P.) 433–439

N-Acetylhexosaminidase, placenta, human, immunochemical characterization of human liver proteins structurally related to (Carroll, M.) 191–196

β-*N*-Acetylhexosaminidase, secretion of isoenzymes of, by cultured human skin fibroblasts (Willcox, P.) 433–439

β-*N*-Acetylhexosaminidases, liver, human, difference in the specificities of, detected by their activities towards oligosaccharides prepared from glycosaminoglycans (Bearpark, T. M. & Stirling, J. L.) 997–1000

Acetylphenylhydrazine, mechanism of the breakdown of human oxyhaemoglobin on reaction with (French, J. K., Winterbourn, C. C. & Carrell, R. W.) 19–26

Acremonium chrysogenum, oxygenation of desacetoxycephalosporin C to desacetylcephalosporin C by 2-oxoglutarate-linked dioxygenases from *Streptomyces clavuligerus* and (Turner, M. K., Farthing, J. E. & Brewer, S. J.) 839–850

Actinidia chinensis, *see* Chinese gooseberry

Actinidin, Chinese-gooseberry, determination of the amino acid sequences of peptides obtained by tryptic digestion of (Carne, A. & Moore, C. H.) 73–83

Actin-like protein, stimulation by 1,25-dihydroxycholecalciferol of the incorporation of leucine into, in chick small-intestine brush border (Wilson, P. W. & Lawson, D. E. M.) 627–631

Adenosine triphosphatase, calcium ion-transporting, sarcoplasmic-reticulum, skeletal-muscle, rabbit, determination of the amino acid sequences of cysteine-containing peptides obtained by tryptic digestion of (Allen, G. & Green, N. M.) 393–402

Adenylate cyclase, corpus luteum, cow, solubilization of, in Lubrol-PX, Triton X-100 or digitonin and the stabilizing effect of sodium fluoride present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924

Adenylate cyclase, time course of and rate-limiting steps in the activation by cholera toxin of, in rat liver (Fischer, J., Kohler, T. R., Lipson, L. G., Flores, J., Witkum, P. A. & Sharp, G. W. G.) 59–64

Albumin, serum, human, removal by transamination and scission of amino acid residues from the peptide representing the cupric ion-transport site of (Sarkar, B., Dixon, H. B. F. & Webster, D.) 895–897

Alcohol dehydrogenase, liver, horse, chemically modified, pH-dependence of intrinsic fluorescence of (Parker, D. M., Hardman, M. J., Plapp, B. V., Holbrook, J. J. & Shore, J. D.) 269–275

Alcohol dehydrogenase, liver, horse, kinetics of the reaction catalysed by, after reductive methylation of its lysine residues (Tsai, C. S.) 483–496

Aldehyde dehydrogenase, potassium ion-activated, baker's-yeast, kinetics and mechanism of the reaction catalysed by (Bostian, K. A. & Betts, G. F.) 787–798

Aldehyde dehydrogenase, potassium ion-activated, baker's-yeast, rapid purification and properties of (Bostian, K. A. & Betts, G. F.) 773–786

Alkanethiols, binding of, and other ligands and spectral shifts in ox heart cytochrome *c* oxidase (Nicholls, P. & Hildebrandt, V.) 65–72

Alloxan-diabetes, *see* Diabetes, alloxan-induced

Amino acid compositions, determination of, and other properties of guinea-pig milk caseins (Craig, R. K., McIlreavy, D. & Hall, R. L.) 633–641

Amino acid sequence, complete, determination of, of a type-I α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of sheep wool α-keratin (Gough, K. H., Inglis, A. S. & Crewther, W. G.) 373–385

Amino acid sequence, complete, determination of, of a type-II α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of sheep wool α-keratin (Crewther, W. G., Inglis, A. S. & McKern, N. M.) 365–371

Amino acid sequence, determination of, of *Porphyra umbilicalis* ferredoxin (Takruri, I., Haslett, B. G., Boulter, D., Andrew, P. W. & Rogers, L. J.) 459–466

Amino acid sequence, determination of, of the major β-chain of rat haemoglobins (Garrick, L. M., Sloan, R. L., Ryan, T. W., Klonowski, T. J. & Garrick, M. D.) 321–330

Amino acid sequence, determination of, of the *N*-terminal 108 amino acid residues of the B chain of subcomponent C1q of component C1 of human complement (Reid, K. B. M. & Thompson, E. O. P.) 863–868

Amino acid sequence, primary, characterization of a region of, of rabbit skeletal-muscle troponin C that is involved in calcium ion-dependent interaction with troponin I (Weeks, R. A. & Perry, S. V.) 449–457

Amino acid sequences, determination of, of cysteine-containing peptides obtained by tryptic digestion of rabbit skeletal-muscle sarcoplasmic-reticulum calcium ion-transporting adenosine triphosphatase (Allen, G. & Green, N. M.) 393–402

Amino acid sequences, determination of, of peptides obtained by tryptic and chymotryptic digestion of a type-II α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of sheep wool α-keratin (Hogg, D. McC., Dowling, L. M. & Crewther, W. G.) 353–363

Amino acid sequences, determination of, of peptides obtained by tryptic digestion of Chinese-gooseberry actinidin (Carne, A. & Moore, C. H.) 73–83

Amino acid sequences, statistical analysis of, of α-helical segments of *S*-carboxymethylkerateine-A from the low-sulphur fraction of sheep wool α-keratin (Elleman, T. C., Crewther, W. G. & Van der Touw, J.) 387–391

α-Amylase, *Bacillus amyloliquefaciens*, role of zinc ions in the stabilization of the dimeric form of (Tellam, R., Winzor, D. J. & Nichol, L. W.) 185–190

α-Amylase, yellow-mealworm, structural studies of monomeric and dimeric wheat-kernel proteins with inhibitory activity towards (Petrucci, T., Sannia, G., Parlamenti, R. & Silano, V.) 229–235

Anti-dinitrophenyl immunoglobulin G2, colostral, cow, thermodynamics of the interaction of ε-dinitrophenyl-L-lysine with (Mukkur, T. K. S.) 39–44

Anti-thrombin III, effect of heparin on the inactivation of ox thrombin by (Machovich, R. & Arányi, P.) 869–875

Aorta, chick-embryo, factors influencing the formation of isodesmosine and desmosine by the action of lysyl oxidase on tropoelastin from (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862

Arginine residue, essential, evidence for, in the recognition site of rabbit skeletal-muscle adenosine 3′:5′-cyclic monophosphate-dependent protein kinase from (Matsuo, M., Huang, C. & Huang, L. C.) 441–447

Arylsulphatase, brain, rat, kinetics and mechanism of the reaction catalysed by (Pennings, E. J. M., Vrielink, R. & Van Kempen, G. M. J.) 299–307

Asialo-carcinoembryonic antigen, effects of the administration of Triton WR1339 and asialo-fetuin on the uptake by mouse liver of carcinoembryonic antigen and, from the blood (Thomas, P. & Jones, M.) 981–983

Asialo-fetuin, effects of the administration of Triton WR1339 and, on the uptake by mouse liver of carcinoembryonic antigen and asialo-carcinoembryonic antigen from the blood (Thomas, P. & Jones, M.) 981–983

Aspartic acid residue 53, structure and enzyme activity of ox pancreatic ribonuclease A after esterification at glutamic acid residue 49 and (Acharya, A. S., Manjula, B. N. & Vithayathil, P. J.) 821–830

Asthma, identification of hyaluronic acid in the pulmonary secretions of human patients with (Sahu, S. & Lynn, W. S.) 565–568

Atolla wyvillei, *see* Jellyfish

Azide, binding of, and other ligands and spectral shifts in ox heart cytochrome *c* oxidase (Nicholls, P. & Hildebrandt, V.) 65–72

Azotobacter chroococcum, electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein component of the nitrogenase complexes of, and other bacteria between +50 and −450mV (O'Donnell, M. J. & Smith, B. E.) 831–838

Azotobacter vinelandii, electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein component of the nitrogenase complexes of, and other bacteria between +50 and −450mV (O'Donnell, M. J. & Smith, B. E.) 831–838

Bacillus amyloliquefaciens, role of zinc ions in the stabilization of the dimeric form of α-amylase from (Tellam, R., Winzor, D. J. & Nichol, L. W.) 185–190

Bacillus megaterium N.C.T.C. 10342, inactivation by phenylglyoxal, butane-2,3-dione and pyridoxal 5′-phosphate of glutamate dehydrogenase and glutamate synthase from (Hemmilä, I. A. & Mäntsälä, P. I.) 53–58

Bacillus megaterium N.C.T.C. 10342, purification and properties of glutamate synthase and glutamate dehydrogenase from (Hemmilä, I. A. & Mäntsälä, P. I.) 45–52

Bacillus polymyxa, electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein component of the nitrogenase complexes of, and other bacteria between +50 and −450mV (O'Donnell, M. J. & Smith, B. E.) 831–838

Bacillus sp., thermophilic, isolation and characterization of isocitrate lyase from (Chell, R. M., Sundaram, T. K. & Wilkinson, A. E.) 165–177

Bile pigments, degradation of the haem residue of abnormal haemoglobins to (Brown, S. B. & Docherty, J. C.) 985–987

Bile, rat, human and dog, inaccuracies in the measurement by ethyl anthranilate diazo and solvent partition methods of conjugated and unconjugated bilirubin in (Ostrow, J. D. & Boonyapisit, S. T.) 263–267

Bilirubin, conjugated and unconjugated, inaccuracies in the measurement by ethyl anthranilate diazo and solvent partition methods of, in rat, human and dog bile (Ostrow, J. D. & Boonyapisit, S. T.) 263–267

Bisulphite, sodium, evidence on the conformation of HeLa-cell 5.8S ribosomal ribonucleic acid from the reaction of, with specific cytidylate residues (Kelly, J. M., Goddard, J. P. & Maden, B. E. H.) 521–532

Blood-group-I activity, association of, with the major human erythrocyte membrane protein (Band 3) (Childs, R. A., Feizi, T., Fukuda, M. & Hakomori, S.-I.) 333–336

Blood-group-Ii activity, isolation and properties of human erythrocyte membrane gangliosides possessing (Feizi, T., Childs, R. A., Hakomori, S.-I. & Powell, M. E.) 245–254

Blood, mouse, effects of the administration of Triton WR1339 and asialo-fetuin on the uptake by the liver of carcinoembryonic antigen and asialo-carcinoembryonic antigen from (Thomas, P. & Jones, M.) 981–983

Blood plasma, human, physical and chemical properties of α_2-macroglobulin from (Hall, P. K. & Roberts, R. C.) 27–38

Bohr effect, alkaline, identification by hydrogen–tritium exchange of the histidine-122α residues of human haemoglobin as groups involved in (Nishikura, K.) 651–657

Brain, ox, solubilization by sodium dodecyl sulphate of purified preparations of myelin from (Crang, A. J. & Rumsby, M. G.) 909–918

Brain, pig, interaction of acetylcholinesterase purified from, with liposomes (Reavill, C. A., Wooster, M. S. & Plummer, D. T.) 851–856

Brain, rat, kinetics and mechanism of the reaction catalysed by arylsulphatase from (Pennings, E. J. M., Vrielink, R. & Van Kempen, G. M. J.) 299–307

Brush-border membrane, small-intestine, rat, purification of maltase/glucoamylase from, and its anomalous dissociation either by heat or by low pH (Flanagan, P. R. & Forstner, G. G.) 553–563

Brush border, small-intestine, chick, stimulation by 1,25-dihydroxycholecalciferol of the incorporation of leucine into an actin-like protein in (Wilson, P. W. & Lawson, D. E. M.) 627–631

Bursa of Fabricius, chicken, comparison of the compositions of glycosphingolipids and gangliosides of, and of chicken, rat and human thymus gland (Narasimhan, R. & Murray, R. K.) 475–482

Butane-2,3-dione, inactivation by phenylglyoxal, pyridoxal 5′-phosphate and, of *Bacillus megaterium* N.C.T.C. 10342 glutamate dehydrogenase and glutamate synthase (Hemmilä, I. A. & Mäntsälä, P. I.) 53–58

Calcium ions, characterization of a region of the primary amino acid sequence of rabbit skeletal-muscle troponin C that is involved in interaction with troponin I dependent on (Weeks, R. A. & Perry, S. V.) 449–457

S-Carboxymethylkerateine-A, wool, sheep, determination of the amino acid sequences of peptides obtained by tryptic and chymotryptic digestion of a type-II α-helical segment of (Hogg, D. McC., Dowling, L. M. & Crewther, W. G.) 353–363

S-Carboxymethylkerateine-A, wool, sheep, determination of the complete amino acid sequence of a type-I α-helical segment of (Gough, K. H., Inglis, A. S. & Crewther, W. G.) 373–385

S-Carboxymethylkerateine-A, wool, sheep, determination of the complete amino acid sequence of a type-II α-helical segment of (Crewther, W. G., Inglis, A. S. & McKern, N. M.) 365–371

S-Carboxymethylkerateine-A, wool, sheep, statistical analysis of the amino acid sequences of α-helical segments of (Elleman, T. C., Crewther, W. G. & Van der Touw, J.) 387–391

Carcinoembryonic antigen, effects of the administration of Triton WR1339 and asialo-fetuin on the uptake by mouse liver of asialo-carcinoembryonic antigen and, from the blood (Thomas, P. & Jones, M.) 981–983

Cartilage, laryngeal, pig, characterization by electric birefringence of proteoglycans from, and of their complexes with hyaluronic acid (Isles, M., Foweraker, A. R., Jennings, B. R., Hardingham, T. & Muir, H.) 237–243

Casein, milk, rabbit, immunochemical characterization of (Al-Sarraj, K., White, D. A. & Mayer, R. J.) 877–883

Caseins, milk, guinea-pig, separation and partial characterization of (Craig, R. K., McIlreavy, D. & Hall, R. L.) 633–641

Catecholamines, biochemical applications of chemically induced nuclear polarization in phenols, peptides and (Muszkat, K. A., Weinstein, M. & Gilon, C.) 993–996

Cells, HeLa, evidence on the conformation of 5.8S ribosomal ribonucleic acid from, from the reaction of sodium bisulphite with specific cytidylate residues (Kelly, J. M., Goddard, J. P. & Maden, B. E. H.) 521–532

Cells, L929, mouse, involvement of deoxyribonucleic acid polymerase β in the biosynthesis of deoxyribonucleic acid in nuclei from (Butt, T. R., Wood, W. M., McKay, E. L. & Adams, R. L. P.) 309–314

Central nervous system, *see* Nervous system, central

Cephalosporin C, oxygenation of desacetoxycephalosporin C to desacetylcephalosporin C by *Acremonium chrysogenum* and *Streptomyces clavuligerus* 2-oxoglutarate-linked dioxygenases and the role of the enzymes in the biosynthesis of (Turner, M. K., Farthing, J. E. & Brewer, S. J.) 839–850

Cervix, cow, structural studies on glycoproteins from mucus secreted by (Roberts, G. P.) 941–947

Chick embryo, factors influencing the formation of isodesmosine and desmosine by the action of lysyl oxidase on tropoelastin from the aorta of (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862

Chick embryo, purification and properties of retinoic acid-binding protein from skin from (Sani, B. P. & Banerjee, C. K.) 643–649

Chick, stimulation by 1,25-dihydroxycholecalciferol of the incorporation of leucine into an actin-like protein in small-intestine brush border of (Wilson, P. W. & Lawson, D. E. M.) 627–631

Chicken, comparison of the compositions of glycosphingolipids and gangliosides of bursa of Fabricius and thymus gland of, and of rat and human thymus gland (Narasimhan, R. & Murray, R. K.) 475–482

Chicken egg, iron-binding properties of ovotransferrin from (Williams, J., Evans, R. W. & Moreton, K.) 533–539

Chicken egg-white, structures of crystals of lysozyme from, and of complexes of the enzyme with *N*-acetylglucosamine and β-methyl *N*-acetylglucosaminide grown in acetate-free medium (Perkins, S. J., Johnson, L. N., Machin, P. A. & Phillips, D. C.) 607–616

Chicken egg-yolk, characterization of glycosphingolipids of (Li, S.-C., Chien, J.-L., Wan, C. C. & Li, Y-T.) 697–699

Chinese gooseberry (*Actinidia chinensis*), determination of the amino acid sequences of peptides obtained by tryptic digestion of actinidin from (Carne, A. & Moore, C. H.) 73–83

Cholecalciferol, 1,25-dihydroxy-, stimulation by, of the incorporation of leucine into an actin-like protein in chick small-intestine brush border (Wilson, P. W. & Lawson, D. E. M.) 627–631

Cholera toxin, time course of and rate-limiting steps in the activation by, of adenylate cyclase in rat liver (Fischer, J., Kohler, T. R., Lipson, L. G., Flores, J., Witkum, P. A. & Sharp, G. W. G.) 59–64

Chondroitin sulphate, interactions between different ox corneal proteoglycans containing keratan sulphate and (Speziale, P., Speziale, M. S., Galligani, L. & Balduini, C.) 935–939

Chromatin, macronuclear, *Tetrahymena pyriformis*, composition of the histones of, containing deoxyribonucleic acid (Jones, R. W.) 155–164

Chromatin, macronuclear, *Tetrahymena pyriformis*, preparation of, containing deoxyribonucleic acid coding for ribosomal ribonucleic acid (Jones, R. W.) 145–153

Chymopapains, use of 2,2′-dipyridyl disulphide in a spectrophotometric method for the detection of, as contaminants in preparations of papain (Baines, B. S. & Brocklehurst, K.) 345–347

Clostridium pasteurianum, electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein component of the nitrogenase complexes of, and other bacteria between +50 and −450mV (O'Donnell, M. J. & Smith, B. E.) 831–838

Cobalophilin, comparison of the properties of intrinsic factor and, present in digestive juices in different parts of the pig gastrointestinal tract (Marcoullis, G., Merivuori, H. & Gräsbeck, R.) 705–712

Collagen, heart-valve, ox, pepsin-solubilized, characterization of (Bashey, R. I., Bashey, H. M. & Jimenez, S. A.) 885–894

Collagen, heart-valve, ox, solubilization of (Jimenez, S. A. & Bashey, R. I.) 337–340

Colon, *see* Intestine, large

Colostrum, cow, thermodynamics of the interaction of ε-dinitrophenyl-L-lysine with anti-dinitrophenyl immunoglobulin G2 from (Mukkur, T. K. S.) 39–44

Complement, human, determination of the amino acid sequence of the *N*-terminal 108 amino acid residues of the B chain of subcomponent C1q of component C1 of (Reid, K. B. M. & Thompson, E. O. P.) 863–868

Component C1, complement, human, determination of the amino acid sequence of the *N*-terminal 108 amino acid residues of the B chain of subcomponent C1q of (Reid, K. B. M. & Thompson, E. O. P.) 863–868

Cornea, ox, interactions between different proteoglycans from (Speziale, P., Speciale, M. S., Galligani, L. & Balduini, C.) 935–939

Corpus luteum, cow, solubilization of adenylate cyclase from, in Lubrol-PX, Triton X-100 or digitonin and the stabilizing effect of sodium fluoride present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924

Creatine kinase, skeletal-muscle, rabbit, properties of matrix-bound dimer and monomer derivatives of, immobilized on Sepharose 4B (Bickerstaff, G. F. & Price, N. C.) 85–93

Cupric ions, removal by transamination and scission of amino acid residues from the peptide representing the transport site for, of human serum albumin (Sarkar, B., Dixon, H. B. F. & Webster, D.) 895–897

Cyanide, binding of, and other ligands and spectral shifts in ox heart cytochrome *c* oxidase (Nicholls, P. & Hildebrandt, V.) 65–72

Cysteine residues, determination of the amino acid sequences of peptides containing, obtained by tryptic digestion of rabbit skeletal-muscle sarcoplasmic-reticulum calcium ion-transporting adenosine triphosphatase (Allen, G. & Green, N. M.) 393–402

Cytidylate residues, specific, evidence on the conformation of HeLa-cell 5.8S ribosomal ribonucleic acid from the reaction of sodium bisulphite with (Kelly, J. M., Goddard, J. P. & Maden, B. E. H.) 521–532

Cytochrome *c* oxidase, heart, ox, binding of ligands and spectral shifts in (Nicholls, P. & Hildebrandt, V.) 65–72

Cytochrome *c* oxidase, heart, ox, magnetic-circular-dichroism studies of haem *a* extracted from, and of its derivatives (Brittain, T., Greenwood, C., Springall, J. P. & Thomson, A. J.) 411–417

Cytochrome *c* oxidase, membrane-bound, mitochondrial, heart, ox, fully reduced, mechanism of reaction of, with oxygen at 176K (Clore, G. M. & Chance, E. M.) 799–810

Cytochrome *c* oxidase, membrane-bound, mitochondrial, heart, ox, mixed-valence-state, ferricyanide-pretreated, mechanism of reaction of, with oxygen at 173K (Clore, G. M. & Chance, E. M.) 811–820

Cytochrome *c*-551 oxidase, *Pseudomonas aeruginosa* N.C.T.C. 6750, oxidation by potassium ferricyanide of (Barber, D., Parr, S. R. & Greenwood, C.) 681–690

Cytochrome *c*-551 oxidase, *Pseudomonas aeruginosa* N.C.T.C. 6750, reaction of oxygen with (Greenwood, C., Barber, D., Parr, S. R., Antonini, E., Brunori, M. & Colosimo, A.) 11–17

Cytosol, heart, pig, comparison of the ionizing groups involved in the binding of reduced nicotinamide-adenine dinucleotide to malate dehydrogenase and lactate dehydrogenase from (Lodola, A., Parker, D. M., Jeck, R. & Holbrook, J. J.) 597–605

Dehydrogenases, nicotinamide–adenine dinucleotide phosphate-dependent, selective retardation of, on chromatography columns of immobilized Procion Red HE-3B (Watson, D. H., Harvey, M. J. & Dean, P. D. G.) 591–596

Deoxycytidine kinase, general method for the purification of, from various mammalian tissues (Baxter, A., Currie, L. M. & Durham, J. P.) 1005–1008

Deoxyribonucleic acid, composition of the histones of *Tetrahymena pyriformis* macronuclear chromatin containing, coding for ribosomal ribonucleic acid (Jones, R. W.) 155–164

Deoxyribonucleic acid, formation and subsequent removal of O^6-methylguanine from, in rat liver and kidney after administration of small doses of dimethylnitrosamine (Pegg, A. E. & Hui, G.) 739–748

Deoxyribonucleic acid, involvement of deoxyribonucleic acid polymerase-β in the biosynthesis of, in mouse L929-cell nuclei (Butt, T. R., Wood, W. M., McKay, E. L. & Adams, R. L. P.) 309–314

Deoxyribonucleic acid polymerase-β, involvement of, in the biosynthesis of deoxyribonucleic acid in mouse L929-cell nuclei (Butt, T. R., Wood, W. M., McKay, E. L. & Adams, R. L. P.) 309–314

Deoxyribonucleic acid, preparation of *Tetrahymena pyriformis* macronuclear chromatin containing, coding for ribosomal ribonucleic acid (Jones, R. W.) 145–153

Deoxyribonucleic acid, thymus-gland, calf, defined molecular-weight distribution of, after extensive sonication (Davis, A. W. & Phillips, D. R.) 179–183

Deoxyribonucleic acids, bifunctional intercalation and sequence specificity in the binding of quinoxaline antibiotics to (Lee, J. S. & Waring, M. J.) 115–128

Deoxyribonucleic acids, changes in mechanism and specificity related to structural changes in the interaction between synthetic analogues of quinoxaline antibiotics and (Lee, J. S. & Waring, M. J.) 129–144

Desacetoxycephalosporin C, oxygenation of, to desacetylcephalosporin C by *Acremonium chrysogenum* and *Streptomyces clavuligerus* 2-oxoglutarate-linked dioxygenases (Turner, M. K., Farthing, J. E. & Brewer, S. J.) 839–850

Desmosine cross-links, structural model for peptides containing, obtained by thermolytic digestion of ox ligamentum nuchae elastin (Mecham, R. P. & Foster, J. A.) 617–625

Desmosine, factors influencing the formation of isodesmosine and, by the action of lysyl oxidase on chick-embryo aorta tropoelastin (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862

Desulfovibrio gigas, determination of the redox potential of the molybdenum-containing iron–sulphur protein from (Moura, J. J. G., Xavier, A. V., Cammack, R., Hall, D. O., Bruschi, M. & Le Gall, J.) 419–425

Diabetes, alloxan-induced, inhibition by starvation and, of the dephosphorylation of inactive pyruvate dehydrogenase complex to the active complex in rat heart mitochondria (Hutson, N. J., Kerbey, A. L., Randle, P. J. & Sugden, P. H.) 669–680

Digitonin, solubilization of cow corpus luteum adenylate cyclase in Lubrol-PX, Triton X-100 or, and the stabilizing effect of sodium fluoride present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924

1,25-Dihydroxycholecalciferol, stimulation by, of the incorporation of leucine into an actin-like protein in chick small-intestine brush border (Wilson, P. W. & Lawson, D. E. M.) 627–631

Dimethylnitrosamine, formation and subsequent removal of O^6-methylguanine from deoxyribonucleic acid in rat liver and kidney after administration of small doses of (Pegg, A. E. & Hui, G.) 739–748

2,4-Dinitrophenylation, inactivation and modification by, of *Mycobacterium smegmatis* lactate oxidase (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 255–262

ε-Dinitrophenyl-L-lysine, thermodynamics of the interaction of, with cow colostral anti-dinitrophenyl immunoglobulin G2 (Mukkur, T. K. S.) 39–44

Dioxygenases, 2-oxoglutarate-linked, *Acremonium chrysogenum* and *Streptomyces clavuligerus*, oxygenation of desacetoxycephalosporin C to desacetylcephalosporin C by (Turner, M. K., Farthing, J. E. & Brewer, S. J.) 839–850

2,2′-Dipyridyl disulphide, use of, in a spectrophotometric method for the detection of chymopapains as contaminants in preparations of papain (Baines, B. S. & Brocklehurst, K.) 345–347

Disulphide bonds, influence of length of the peptide chain on the formation of, in neurohypophysial hormones and their analogues (Moore, G.) 403–409

6,6′-Dithiodinicotinate, half-of-the-sites oxidation by, of ox liver uridine diphosphate glucose dehydrogenase (Franzen, J. S., Marchetti, P., Ishman, R., Ashcom, J. & Feingold, D. S.) 701–704

Dodecyl sulphate, sodium, solubilization by, of purified preparations of ox central-nervous-system myelin (Crang, A. J. & Rumsby, M. G.) 909–917

Egg, chicken, iron-binding properties of ovotransferrin from (Williams, J., Evans, R. W. & Moreton, K.) 535–541

Egg-white, chicken, structures of crystals of lysozyme from, and of complexes of the enzyme with *N*-acetylglucosamine and β-methyl *N*-acetylglucosaminide grown in acetate-free medium (Perkins, S. J., Johnson, L. N., Machin, P. A. & Phillips, D. C.) 607–616

Egg-yolk, chicken, characterization of glycosphingolipids of (Li, S.-C., Chien, J.-L., Wan, C. C. & Li, Y.-T.) 697–699

Elastin, factors influencing the formation of isodesmosine and desmosine by the action of lysyl oxidase on chick-embryo aorta tropoelastin and their relevance to the biosynthesis of (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862

Elastin, ligamentum nuchae, ox, structural model for peptides containing desmosine cross-links obtained by thermolytic digestion of (Mecham, R. P. & Foster, J. A.) 617–625

Enzymes, diffusional increase and decrease of Michaelis constants for two-substrate reactions catalysed by (Engasser, J.-M. & Hisland, P.) 341–343

Erythrocytes, human, association of blood-group-I activity with the major membrane protein (Band 3) of (Childs, R. A., Feizi, T., Fukuda, M. & Hakomori, S.-I.) 333–336

Erythrocytes, human, isolation and properties of blood-group-Ii-active gangliosides from the membrane of (Feizi, T., Childs, R. A., Hakomori, S.-I. & Powell, M. E.) 245–254

Erythrocytes, human, purification and properties of γ-glutamylcyclotransferase from (Board, P. G., Moore, K. A. & Smith, J. E.) 427–431

Erythrocytes, ox, similarity of the two high-molecular-weight polypeptides of spectrin from the membrane of (Dunn, M. J., Kemp, R. B. & Maddy, A. H.) 197–205

Erythrocytes, pigeon, composition of the protein and lipid components of the membrane of (Watts, C. & Wheeler, K. P.) 899–907

Escherichia coli, molecular weights and the problem of aggregation of enzymes involved in the biosynthesis of proline in (Hayzer, D. J. & Moses, V.) 219–228

Escherichia coli, potential errors arising from the use of a bioradiological procedure for assay of the biosynthesis of proline by cell-free extracts of (Hayzer, D. J. & Moses, V.) 207–217

Ethylene, electron-paramagnetic-resonance evidence for the formation of transient complexes of acetylene and, with *Klebsiella pneumoniae* nitrogenase complex (Lowe, D. J., Eady, R. R. & Thorneley, R. N. F.) 277–290

Euglena gracilis, purification and immunochemical characterization of malate synthase from (Woodcock, E. & Merrett, M. J.) 95–101

Eye, ox, interactions between different corneal proteoglycans from (Speziale, P., Speziale, M. S., Galligani, L. & Balduini, C.) 935–939

Fatty acid synthetase, mammary-gland, rabbit, lactating, molecular weight and subunit size of (Grunnet, I. & Knudsen, J.) 929–933

Ferredoxin, *Porphyra umbilicalis*, determination of the amino acid sequence of (Takruri, I., Haslett, B G., Boulter, D., Andrew, P. W. & Rogers, L. J.) 459–466

Ferricyanide, potassium, oxidation by, of *Pseudomonas aeruginosa* N.C.T.C. 6750 cytochrome *c*-551 oxidase (Barber, D., Parr, S. R. & Greenwood, C.) 681–690
Ferritins, human, properties of isomers of, from various tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977
Fibroblasts, skin, human, cultured, secretion of isoenzymes of β-*N*-acetylglucosaminidase by (Willcox, P.) 433–439
Fibroblasts, synovial, rabbit, properties of proteinases from, and polymorphonuclear leucocytes with activity towards proteoglycans (Werb, Z., Dingle, J. T., Reynolds, J. J. & Barrett, A. J.) 949–958
Fluorescence, intrinsic, pH-dependent changes of, of chemically modified horse liver alcohol dehydrogenase (Parker, D. M., Hardman, M. J., Plapp, B. V., Holbrook, J. J. & Shore, J. D.) 269–275
Fluoride, sodium, solubilization of cow corpus luteum adenylate cyclase in Lubrol-PX, Triton X-100 or digitonin and the stabilizing effect of, present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924
1-Fluoro-2,4-dinitrobenzene, inactivation and modification by, of *Mycobacterium smegmatis* lactate oxidase (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 255–262
α-L-Fucosidase, liver, human, anomalous gel-filtration behaviour of, and other sialoglycoproteins (Alhadeff, J. A.) 315–319

Gangliosides, blood-group-Ii-active, membrane, erythrocyte, human, isolation and properties of (Feizi, T., Childs, R. A., Hakomori, S.-I. & Powell, M. E.) 245–254
Gangliosides, comparison of the compositions of glycosphingolipids and, of chicken bursa of Fabricius and of chicken, rat and human thymus gland (Narasimhan, R. & Murray, R. K.) 475–482
Gastrointestinal tract, pig, comparison of the properties of intrinsic factor and cobalophilin present in digestive juices in different parts of (Marcoullis, G., Merivuori, H. & Gräsbeck, R.) 705–712
Glucoamylase/maltase, brush-border-membrane, small-intestine, rat, purification of, and its anomalous dissociation either by heat or by low pH (Flanagan, P. R. & Forstner, G. G.) 553–563
Glutamate dehydrogenase, *Bacillus megaterium* N.C.T.C. 10342, inactivated by phenylglyoxal, butane-2,3-dione and pyridoxal 5′-phosphate of, and glutamate synthase (Hemmilä, I. A. & Mäntsälä, P. I.) 53–58
Glutamate dehydrogenase, *Bacillus megaterium* N.C.T.C. 10342, purification and properties of, and glutamate synthase (Hemmilä, I. A. & Mäntsälä, P. I.) 45–52
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-dependent, *Neurospora crassa*, use of affinity chromatography on immobilized Procion Red HE-3B in the purification of (Watson, D. H., Harvey, M. J. & Dean, P. D. G.) 591–596
Glutamate synthase, *Bacillus megaterium* N.C.T.C. 10342, inactivation by phenylglyoxal, butane-2,3-dione and pyridoxal 5′-phosphate of, and glutamate dehydrogenase (Hemmilä, I. A. & Mantsälä, P. I.) 53–58
Glutamate synthase, *Bacillus megaterium* N.C.T.C. 10342, purification and properties of, and glutamate dehydrogenase (Hemmilä, I. A. & Mäntsälä, P. I.) 45–52
Glutamic acid residue 49, structure and enzyme activity of ox pancreatic ribonuclease A after esterification of aspartic acid residue 53 and (Acharya, A. S., Manjula, B. N. & Vithayathil, P. J.) 821–830
Glutamyl kinases, *Escherichia coli*, molecular weights of, and other enzymes involved in the biosynthesis of proline and the problem of enzyme aggregation (Hayzer, D. J. & Moses, V.) 219–228
γ-Glutamyl phosphate reductase, *Escherichia coli*, molecular weights of, and other enzymes involved in the biosynthesis of proline and the problem of enzyme aggregation (Hayzer, D. J. & Moses, V.) 219–228
γ-Glutamylcyclotransferase, erythrocyte, human, purification and properties of (Board, P. G., Moore, K. A. & Smith, J. E.) 427–431
Glutathione, role of γ-glutamylcyclotransferase in the biosynthesis of, by human erythrocytes (Board, P. G., Moore, K. A. & Smith, J. E.) 427–431
rac-Glycerol 1:2-cyclic phosphate phosphodiesterase, kidney-cortex, pig, purification and properties of (Clarke, N. & Dawson, R. M. C.) 579–589
Glycoprotein, high-molecular-weight, mucus, large-intestine, pig, isolation and characterization of (Marshall, T. & Allen, A.) 569–578
Glycoproteins, mucus, cervical, cow, structural studies on (Roberts, G. P.) 941–947
Glycoproteins, sialo-, anomalous gel-filtration behaviour of (Alhadeff, J. A.) 315–319
Glycosaminoglycans, difference in the specificities of human liver β-*N*-acetylhexosaminidases detected by their activities towards oligosaccharides prepared from (Bearpark, T. M. & Stirling, J. L.) 997–1000
Glycosaminoglycans, oxidation by periodate and the configurations of, in solution (Scott, J. E. & Tigwell, M. J.) 103–114
Glycosphingolipids, comparison of the compositions of gangliosides and, of chicken bursa of Fabricius and of chicken, rat and human thymus gland (Narasimhan, R. & Murray, R. K.) 475–482
Glycosphingolipids, egg-yolk, chicken, characterization of (Li, S.-C., Chien, J.-L., Wan, C. C. & Li, Y.-T.) 697–699
Glyoxylic acid cycle, isolation and characterization of isocitrate lyase and its role in, in a thermophilic *Bacillus* sp. (Chell, R. M., Sundaram, T. K. & Wilkinson, A. F.) 165–177
Granulocytes, *see* Leucocytes, polymorphonuclear

Haem *a*, magnetic-circular-dichroism studies of, extracted from ox heart cytochrome *c* oxidase and of its derivatives (Brittain, T., Greenwood, C., Springall, J. P. & Thomson, A. J.) 411–417
Haem residue, degradation of, of abnormal human haemoglobins (Brown, S. B. & Docherty, J. C.) 985–987
Haemoglobin, human, identification by hydrogen–tritium exchange of the histidine-122α residues of, as groups involved in the alkaline Bohr effect (Nishikura, K.) 651–657

Haemoglobin, oxy-, human, mechanism of the breakdown of, on reaction with acetylphenylhydrazine (French, J. K., Winterbourn, C. C. & Carrell, R. W.) 19–26

Haemoglobins, human, abnormal, degradation of the haem residue of (Brown, S. B. & Docherty, J. C.) 985–987

Haemoglobins, rat, determination of the amino acid sequence of the major β-chain of (Garrick, L. M., Sloan, R. L., Ryan, T. W., Klonowski, T. J. & Garrick, M. D.) 321–330

Heart, human, properties of isoferritins from, and other tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977

Heart, ox, binding of ligands and spectral shifts in cytochrome *c* oxidase from (Nicholls, P. & Hildebrandt, V.) 65–72

Heart, ox, magnetic-circular-dichroism studies of haem *a* extracted from cytochrome *c* oxidase from, and of its derivatives (Brittain, T., Greenwood, C., Springall, J. P. & Thomson, A. J.) 411–417

Heart, ox, mechanism of reaction of ferricyanide-pretreated mixed-valence-state mitochondrial membrane-bound cytochrome *c* oxidase from, with oxygen at 173K (Clore, G. M. & Chance, E. M.) 811–820

Heart, ox, mechanism of reaction of fully reduced mitochondrial membrane-bound cytochrome *c* oxidase from, with oxygen at 176K (Clore, G. M. & Chance, E. M.) 799–810

Heart, pig, comparison of the ionizing groups involved in the binding of reduced nicotinamide–adenine dinucleotide to cytosolic malate dehydrogenase and lactate dehydrogenase from (Lodola, A., Parker, D. M., Jeck, R. & Holbrook, J. J.) 597–605

Heart, pig, subunit stoicheiometry in the regulation by phosphorylation of the activity of pyruvate dehydrogenase complex from (Sugden, P. H. & Randle, P. J.) 659–668

Heart, pig, use of the sulphite adduct of oxidized nicotinamide–adenine dinucleotide to study ionizations of and kinetics of the reactions catalysed by lactate dehydrogenase and malate dehydrogenase from (Parker, D. M., Lodola, A. & Holbrook, J. J.) 959–967

Heart, rat, inhibition by alloxan-diabetes and starvation of the dephosphorylation of inactive pyruvate dehydrogenase complex to the active complex in mitochondria from (Hutson, N. J., Kerbey, A. L., Randle, P. J. & Sugden, P. H.) 669–680

Heart valves, ox, characterization of pepsin-solubilized collagen from (Bashey, R. I., Bashey, H. M. & Jimenez, S. A.) 885–894

Heart valves, ox, solubilization of collagen from (Jimenez, S. A. & Bashey, R. I.) 337–340

HeLa cells, evidence on the conformation of 5.8S ribosomal ribonucleic acid from, from the reaction of sodium bisulphite with specific cytidylate residues (Kelly, J. M., Goddard, J. P. & Maden, B. E. H.) 521–532

Hen, *see* Chicken

Heparin, effect of, on the inactivation of ox thrombin by anti-thrombin III (Machovich, R. & Arányi, P.) 869–875

Heparin, retention at low temperatures of rabbit reticulocyte ribosomes on, bound to Sepharose 4B (Hradec, J. & Kříž, O.) 349–352

Hexosaminidase, *see N*-Acetylhexosaminidase

Histidine-122α residues, identification by hydrogen–tritium exchange of, of human haemoglobin as groups involved in the alkaline Bohr effect (Nishikura, K.) 651–657

Histidine residues, involvement of, in the binding of reduced nicotinamide–adenine dinucleotide to pig heart cytosolic malate dehydrogenase and lactate dehydrogenase (Lodola, A., Parker, D. M., Jeck, R. & Holbrook, J. J.) 597–605

Histones, composition of, of *Tetrahymena pyriformis* macronuclear chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid (Jones, R. W.) 155–164

Hyaluronic acid, characterization by electric birefringence of pig laryngeal cartilage proteoglycans and of their complexes with (Isles, M., Fowerаker, A. R., Jennings, B. R., Hardingham, T. & Muir, H.) 237–243

Hyaluronic acid, identification of, in the pulmonary secretions of human patients with asthma (Sahu, S. & Lynn, W. S.) 565–568

Hyaluronic acid, oxidation by periodate and the configurations of, and other glycosaminoglycans in solution (Scott, J. E. & Tigwell, M. J.) 103–114

4-Hydroxy-3-methoxyphenylethylene glycol sulphate, kinetics and mechanism of the reaction catalysed by rat brain arylsulphatase leading to the formation of (Pennings, E. J. M., Vrielink, R. & Van Kempen, G. M. J.) 299–307

3-Hydroxy-3-methylglutaryl-coenzyme A synthase, mitochondrial, liver, ox, purification and properties of (Page, M. A. & Tubbs, P. K.) 925–928

Iduronate residues, oxidation by periodate and the conformation of, of glycosaminoglycans in solution (Scott, J. E. & Tigwell, M. J.) 103–114

Immunoglobulin A (myeloma protein), role of nitro groups in the binding of nitroaromatic compounds to, from mouse plasma-cell tumour MOPC 315 cells (Gettins, P., Givol, D. & Dwek, R. A.) 713–722

Immunoglobulin G2, anti-dinitrophenyl, colostral, cow, thermodynamics of the interaction of ε-dinitrophenyl-L-lysine with (Mukkur, T. K. S.) 39–44

Intestine, large, pig, isolation and characterization of the high-molecular-weight glycoprotein from mucus secreted by (Marshall, T. & Allen, A.) 569–578

Intestine, small, chick, stimulation by 1,25-dihydroxycholecalciferol of the incorporation of leucine into an actin-like protein in brush border of (Wilson, P. W. & Lawson, D. E. M.) 627–631

Intestine, small, pig, comparison of the properties of intrinsic factor and cobalophilin present in digestive juices present in, and stomach (Marcoullis, G., Merivuori, H. & Gräsbeck, R.) 705–712

Intestine, small, rat, inactivation of native enzymes by a neutral proteinase from smooth muscle of (Beynon, R. J. & Kay, J.) 291–298

Intestine, small, rat, purification of brush-border membrane maltase/glucoamylase from, and its anomalous dissociation either by heat or by low pH (Flanagan, P. R. & Forstner, G. G.) 553–563

Intrinsic factor, comparison of the properties of cobalophilin and, present in digestive juices in different parts of the pig gastrointestinal tract (Marcoullis, G., Merivuori, H. & Gräsbeck, R.) 705–712
Ionic strength, effects of pH and, on the allosteric interactions of ox pancreatic ribonuclease with substrates and substrate analogues (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 1–4
Iron, binding of, by chicken-egg ovotransferrin (Williams, J., Evans, R. W. & Moreton, K.) 533–539
Iron, binding of different donors of, to human serum transferrin and isolation of iron-binding fragments from the *N*-terminal and *C*-terminal regions of the protein (Evans, R. W. & Williams, J.) 543–552
Iron, content of, of isoferritins from various human tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977
Iron, derivation of equations relating the fractional saturation of *N*-terminal and *C*-terminal sites to the number of atoms of, bound per molecule of chicken-egg ovotransferrin (Milne, J. G. C.) 542
Iron–sulphur protein, molybdenum-containing, *Desulfovibrio gigas*, determination of the redox potential of (Moura, J. J. G., Xavier, A. V., Cammack, R., Hall, D. O., Bruschi, M. & Le Gall, J.) 419–425
Isocitrate lyase, *Bacillus* sp., thermophilic, isolation and characterization of (Chell, R. M., Sundaram, T. K. & Wilkinson, A. E.) 165–177
Isodesmosine, factors influencing the formation of desmosine and, by the action of lysyl oxidase on chick-embryo aorta tropoelastin (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862
Isoferritins, human, properties of, from various tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977

Jellyfish (*Atolla wyvillei* and *Periphylla periphylla*), use of high-pressure liquid chromatography with solvent systems containing phase-transfer agents for the direct determination of porphyrin carboxylic acids in, and other biological samples (Bonnett, R., Charalambides, A. A., Jones, K., Magnus, I. A. & Ridge, R. J.) 693–696

Keratan sulphate, interactions between different ox corneal proteoglycans containing chondroitin sulphate and (Speziale, P., Speziale, M. S., Galligani, L. & Balduini, C.) 935–939
α-Keratin, wool, sheep, determination of the amino acid sequences of peptides obtained by tryptic and chymotryptic digestion of a type-II α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of (Hogg, D. McC., Dowling, L. M. & Crewther, W. G.) 353–363
α-Keratin, wool, sheep, determination of the complete amino acid sequence of a type-I α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of (Gough, K. H., Inglis, A. S. & Crewther, W. G.) 373–385
α-Keratin, wool, sheep, determination of the complete amino acid sequence of a type-II α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of (Crewther, W. G., Inglis, A. S. & McKern, N. M.) 365–371
α-Keratin, wool, sheep, statistical analysis of the amino acid sequences of α-helical segments of *S*-carboxymethylkerateine-A from the low-sulphur fraction of (Elleman, T. C., Crewther, W. G. & Van der Touw, J.) 387–391
Kidney, calf, lack of tissue specificity of the high-mobility group of chromosomal non-histone proteins from, and other tissues (Rabbani, A., Goodwin, G. H. & Johns, E. W.) 497–505
Kidney cortex, pig, purification and properties of *rac*-glycerol 1:2-cyclic phosphate phosphodiesterase from (Clarke, N. & Dawson, R. M. C.) 579–589
Kidney, human, properties of isoferritins from, and other tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977
Kidney, rat, formation and subsequent removal of O^6-methylguanine from deoxyribonucleic acid in, and liver after administration of small doses of dimethylnitrosamine (Pegg, A. E. & Hui, G.) 739–748
Kidney, sheep, purification and properties of 3-oxo acid coenzyme A-transferase from (Sharp, J. A. & Edwards, M. R.) 759–765
Klebsiella pneumoniae, electron-paramagnetic-resonance evidence for the formation of transient complexes of acetylene and ethylene with nitrogenase complex from (Lowe, D. J., Eady, R. R. & Thorneley, R. N. F.) 277–290
Klebsiella pneumoniae, electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein component of the nitrogenase complexes of, and other bacteria between +50 and −450 mV (O'Donnell, M. J. & Smith, B. E.) 831–838

L929 cells, mouse, involvement of deoxyribonucleic acid polymerase β in the biosynthesis of deoxyribonucleic acid in nuclei from (Butt, T. R., Wood, W. M., McKay, E. L. & Adams, R. L. P.) 309–314
Lactate dehydrogenase, cytosolic, heart, pig, comparison of the ionizing groups involved in the binding of reduced nicotinamide–adenine dinucleotide to, and malate dehydrogenase (Lodola, A., Parker, D. M., Jeck, R. & Holbrook, J. J.) 597–605
Lactate dehydrogenase, heart, pig, use of the sulphite adduct of oxidized nicotinamide–adenine dinucleotide to study ionizations of and kinetics of the reactions catalysed by, and malate dehydrogenase (Parker, D. M., Lodola, A. & Holbrook, J. J.) 959–967
Lactate dehydrogenase, seminal-fluid, human, use of affinity chromatography for the purification of isoenzyme X of (Kolk, A. H. J., Van Kuyk, L. & Boettcher, B.) 767–771
Lactate oxidase, *Mycobacterium smegmatis*, inactivation and modification by 1-fluoro-2,4-dinitrobenzene of (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 255–262
Lactation, immunochemical characterization of casein secreted by rabbit mammary gland during (Al-Sarraj, K., White, D. A. & Mayer, R. J.) 877–883
Lactation, molecular weight and subunit size of rabbit mammary-gland fatty acid synthetase obtained during (Grunnet, I. & Knudsen, J.) 929–933
Large intestine, *see* Intestine, large

Larynx, pig, characterization by electric birefringence of proteoglycans from cartilage of, and of their complexes with hyaluronic acid (Isles, M., Foweraker, A. R., Jennings, B. R., Hardingham, T. & Muir, H.) 237–243

Leucine, stimulation by 1,25-dihydroxycholecalciferol of the incorporation of, into an actin-like protein in chick small-intestine brush border (Wilson, P. W. & Lawson, D. E. M.) 627–631

Leucocytes, polymorphonuclear, rabbit, properties of proteinases from, and synovial fibroblasts with activity towards proteoglycans (Werb, Z., Dingle, J. T., Reynolds, J. J. & Barrett, A. J.) 949–958

Ligamentum nuchae, ox, structural model for peptides containing desmosine cross-links obtained by thermolytic digestin of elastin from (Mecham, R. P. & Foster, J. A.) 617–625

Lipids, composition of the proteins and, present in the pigeon erythrocyte membrane (Watts, C. & Wheeler, K. P.) 899–907

Lipoproteins, serum, rainbow-trout, isolation and partial characterization of, and apolipoproteins (Skinner, E. R. & Rogie, A.) 507–520

Liposomes, interaction of purified pig brain acetylcholinesterase with (Reavill, C. A., Wooster, M. S. & Plummer, D. T.) 851–856

Liver, calf, lack of tissue specificity of the high-mobility group of chromosomal non-histone proteins from, and other tissues (Rabbani, A., Goodwin, G. H. & Johns, E. W.) 497–505

Liver, horse, kinetics of the reaction catalysed by alcohol dehydrogenase from, after reductive methylation of its lysine residues (Tsai, C. S.) 483–496

Liver, horse, pH-dependent changes of intrinsic fluorescence of chemically modified alcohol dehydrogenase from (Parker, D. M., Hardman, M. J., Plapp, B. V., Holbrook, J. J. & Shore, J. D.) 269–275

Liver, human, anomalous gel-filtration behaviour of α-L-fucosidase from, and other sialoglycoproteins (Alhadeff, J. A.) 315–319

Liver, human, difference in the specificities of β-*N*-acetylhexosaminidases from, detected by their activities towards oligosaccharides prepared from glycosaminoglycans (Bearpark, T. M. & Stirling, J. L.) 997–1000

Liver, human, immunochemical characterization of proteins from, that are structurally related to human placenta *N*-acetyl-β-D-glucosaminidase (Carroll, M.) 191–196

Liver, human, properties of isoferritins from, and other tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977

Liver, mouse, effects of the administration of Triton WR1339 and asialo-fetuin on the uptake by, of carcinoembryonic antigen and asialo-carcinoembryonic antigen from the blood (Thomas, P. & Jones, M.) 981–983

Liver, ox, half-of-the-sites oxidation by 6,6′-dithiodinicotinate of uridine diphosphate glucose dehydrogenase from (Franzen, J. S., Marchetti, P., Ishman, R., Ashcom, J. & Feingold, D. S.) 701–704

Liver, ox, purification and properties of mitochondrial 3-hydroxy-3-methylglutaryl-coenzyme A synthase from (Page, M. A. & Tubbs, P. K.) 925–928

Liver, rat, formation and subsequent removal of O^6-methylguanine from deoxyribonucleic acid in, and kidney after administration of small doses of dimethylnitrosamine (Pegg, A. E. & Hui, G.) 739–748

Liver, rat, substrate specificity and properties of uridine diphosphate glucuronyltransferase purified to apparent homogeneity from, of animals pretreated with phenobarbital (Burchell, B.) 749–757

Liver, rat, time course of and rate-limiting steps in the activation by cholera toxin of adenylate cyclase in (Fischer, J., Kohler, T. R., Lipson, L. G., Flores, J., Witkum, P. A. & Sharp, G. W. G.) 59–64

Lubrol-PX, solubilization of cow corpus luteum adenylate cyclase in Triton X-100, digitonin or, and the stabilizing effect of sodium fluoride present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924

Lung, human, identification of hyaluronic acid in the pulmonary secretions of, of patients with asthma (Sahu, S. & Lynn, W. S.) 565–568

Lysine residues, effect of carbamoylation of, on the allosteric interactions of ox pancreatic ribonuclease with substrates and substrate analogues (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 1–4

Lysine residues, kinetics of the reaction catalysed by horse liver alcohol dehydrogenase after reductive methylation of (Tsai, C. S.) 483–496

Lysosomes, skin-fibroblast, human, cultured, secretion of isoenzymes of β-*N*-acetylglucosaminidase by (Willcox, P.) 433–439

Lysozyme, egg-white, chicken, structures of crystals of, and of complexes of the enzyme with *N*-acetylglucosamine and β-methyl *N*-acetylglucosaminide grown in acetate-free medium (Perkins, S. J., Johnson, L. N., Machin, P. A. & Phillips, D. C.) 607–616

Lysyl oxidase, cartilage, chick-embryo, factors influencing the formation of isodesmosine and desmosine by the action of, on chick-embryo aorta tropoelastin (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862

α_2-Macroglobulin, plasma, human, physical and chemical properties of (Hall, P. K. & Roberts, R. C.) 27–38

Macronuclei, *Tetrahymena pyriformis*, composition of the histones of chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid from (Jones, R. W.) 155–164

Macronuclei, *Tetrahymena pyriformis*, preparation of chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid from (Jones, R. W.) 145–153

Malate dehydrogenase, cytosolic, heart, pig, comparison of the ionizing groups involved in the binding of reduced nicotinamide–adenine dinucleotide to, and lactate dehydrogenase (Lodola, A., Parker, D. M., Jeck, R. & Holbrook, J. J.) 597–605

Malate dehydrogenase, heart, pig, use of the sulphite adduct of oxidized nicotinamide–adenine dinucleotide to study ionizations of and kinetics of the reactions catalysed by, and lactate dehydrogenase (Parker, D. M., Lodola, A. & Holbrook, J. J.) 959–967

Malate synthase, *Euglena gracilis*, purification and immunochemical characterization of (Woodcock, E. & Merrett, M. J.) 95–101

Maltase/glucoamylase, brush-border-membrane, small-intestine, rat, purification of, and its anomalous dissociation either by heat or by low pH (Flanagan, P. R. & Forstner, G. G.) 553–563
Mammary gland, rabbit lactating, immunochemical characterization of casein secreted by (Al-Sarraj, K., White, D. A. & Mayer, R. J.) 879–885
Mammary gland, rabbit, lactating, molecular weight and subunit size of fatty acid synthetase from (Grunnet, I. & Knudsen, J.) 929–933
Mealworm, yellow (*Tenebrio molitor*), structural studies of monomeric and dimeric wheat-kernel proteins with inhibitory activity towards α-amylase from (Petrucci, T., Sannia, G., Parlamenti, R. & Silano, V.) 229–235
Membrane, brush-border, small-intestine, rat, purification of maltase/glucoamylase from, and its anomalous dissociation either by heat or by low pH (Flanagan, P. R. & Forstner, G. G.) 553–563
Membrane, erythrocyte, human, isolation and properties of blood-group-Ii-active gangliosides from (Feizi, T., Childs, R. A., Hakomori, S.-I. & Powell, M. E.) 235–254
Membrane, erythrocyte, ox, similarity of the two high-molecular-weight polypeptides of spectrin from (Dunn, M. J., Kemp, R. B. & Maddy, A. H.) 197–205
Membrane, erythrocyte, pigeon, composition of the protein and lipid components of (Watts, C. & Wheeler, K. P.) 899–907
Membrane, mitochondrial, heart, ox, mechanism of reaction of ferricyanide-pretreated mixed-valence-state cytochrome *c* oxidase bound to, with oxygen at 173K (Clore, G. M. & Chance, E. M.) 811–820
Membrane, mitochondrial, heart, ox, mechanism of reaction of fully reduced cytochrome *c* oxidase bound to, with oxygen at 176K (Clore, G. M. & Chance, E. M.) 799–810
Membrane protein (Band 3), erythrocyte, human, major, association of blood-group-I activity with (Childs, R. A., Feizi, T., Fukuda, M. & Hakomori, S.-I.) 333–336
β-Methyl *N*-acetylglucosaminide, structures of crystals of chicken egg-white lysozyme and of complexes of the enzyme with *N*-acetylglucosamine and, grown in acetate-free medium (Perkins, S. J., Johnson, L. N., Machin, P. A. & Phillips, D. C.) 607–616
Methylation, reductive, effect of, of lysine residues on the kinetics of the reaction catalysed by horse liver alcohol dehydrogenase (Tsai, C. S.) 483–496
O^6-Methylguanine, formation and subsequent removal of, from deoxyribonucleic acid in rat liver and kidney after administration of small doses of dimethylnitrosamine (Pegg, A. E. & Hui, G.) 739–748
Michaelis constants, diffusional increase and decrease of, for two-substrate reactions catalysed by enzymes (Engasser, J.-M. & Hisland, P.) 341–343
Milk, guinea-pig, separation and partial characterization of caseins from (Craig, R. K., McIlreavy, D. & Hall, R. L.) 633–641
Milk, rabbit, immunochemical characterization of casein of (Al-Sarraj, K., White, D. A. & Mayer, R. J.) 877–883
Mitochondria, heart, ox, mechanism of reaction of ferricyanide-pretreated mixed-valence-state membrane-bound cytochrome *c* oxidase from, with oxygen at 173K (Clore, G. M. & Chance, E. M.) 811–820
Mitochondria, heart, ox, mechanism of reaction of fully reduced membrane-bound cytochrome *c* oxidase from, with oxygen at 176K (Clore, G. M. & Chance, E. M.) 799–810
Mitochondria, heart, rat, inhibition by alloxan-diabetes and starvation of the dephosphorylation of inactive pyruvate dehydrogenase complex to the active complex in (Hutson, N. J., Kerbey, A. L., Randle, P. J. & Sugden, P. H.) 669–680
Mitochondria, liver, ox, purification and properties of 3-hydroxy-3-methylglutaryl-coenzyme A synthase from (Page, M. A. & Tubbs, P. K.) 925–928
Molybdenum, determination of the redox potential of the *Desulfovibrio gigas* iron–sulphur protein containing (Moura, J. J. G., Xavier, A. V., Cammack, R., Hall, D. O., Bruschi, M. & Le Gall, J.) 419–425
Molybdenum–iron protein component, electron-paramagnetic-resonance studies on the redox properties of, of various bacterial nitrogenase complexes between +50 and −450mV (O'Donnell, M. J. & Smith, B. E.) 831–838
Mucus, cervical, cow, structural studies on glycoproteins from (Roberts, G. P.) 941–947
Mucus, large-intestine, pig, isolation and characterization of the high-molecular-weight glycoprotein from (Marshall, T. & Allen, A.) 569–578
Muscle, skeletal, rabbit, characterization of a region of the primary amino acid sequence of troponin C from, that is involved in calcium ion-dependent interaction with troponin I (Weeks, R. A. & Perry, S. V.) 449–457
Muscle, skeletal, rabbit, determination of the amino acid sequences of cysteine-containing peptides obtained by tryptic digestion of sarcoplasmic-reticulum calcium ion-transporting adenosine triphosphatase from (Allen, G. & Green, N. M.) 393–402
Muscle, skeletal, rabbit, evidence for an essential arginine residue in the recognition site of adenosine 3′:5′-cyclic monophosphate-dependent protein kinase from (Matsuo, M., Huang, C. & Huang, L. C.) 441–447
Muscle, skeletal, rabbit, properties of matrix-bound dimer and monomer derivatives of immobilized creatine kinase from (Bickerstaff, G. F. & Price, N. C.) 85–93
Muscle, smooth, small-intestinal, rat, inactivation of native enzymes by a neutral proteinase from (Beynon, R. J. & Kay, J.) 291–298
Mycobacterium smegmatis, inactivation and modification by 1-fluoro-2,4-dinitrobenzene of lactate oxidase from (Choong, Y. S., Shepherd, M. G. & Sullivan, P. A.) 255–262
Myelin, central-nervous-system, ox, solubilization by sodium dodecyl sulphate of purified preparations of (Crang, A. J. & Rumsby, M. G.) 909–917
Myelin, sciatic-nerve, mouse, composition of the proteins of, from Quaking and Jimpy mutants (Matthieu, J.-M.) 989–991
Myeloma protein (immunoglobulin A), role of nitro groups in the binding of nitroaromatic compounds to, from mouse plasma-cell tumour MOPC 315 cells (Gettins, P., Givol, D. & Dwek, R. A.) 713–722

Nerve, sciatic, mouse, composition of the proteins of the myelin of, from Quaking and Jimpy mutants (Matthieu, J.-M.) 989–991

Nervous system, central, ox, solubilization by sodium dodecyl sulphate of purified preparations of myelin from (Crang, A. J. & Rumsby, M. G.) 909–917

Neurohypophysis, influence of length of the peptide chain on the formation of disulphide bonds in hormones biosynthesized by, and their analogues (Moore, G.) 403–409

Neurospora crassa, use of affinity chromatography on immobilized Procion Red HE-3B in the purification of nicotinamide–adenine dinucleotide phosphate-dependent glutamate dehydrogenase from (Watson, D. H., Harvey, M. J. & Dean, P. D. G.) 591–596

Neutral proteinase, *see* Proteinase, neutral

Nicotinamide–adenine dinucleotide, oxidized, use of the sulphite adduct of, to study ionizations of and kinetics of the reactions catalysed by pig heart lactate dehydrogenase and malate dehydrogenase (Parker, D. M., Lodola, A. & Holbrook, J. J.) 959–967

Nicotinamide–adenine dinucleotide phosphate, selective retardation of dehydrogenases dependent on, on chromatography columns of immobilized Procion Red HE-3B (Watson, D. H., Harvey, M. J. & Dean, P. D. G.) 591–596

Nicotinamide–adenine dinucleotide, reduced, comparison of the ionizing groups involved in the binding of, to pig heart cytosolic malate dehydrogenase and lactate dehydrogenase (Lodola, A., Parker, D. M., Jeck, R. & Holbrook, J. J.) 597–605

Nitro groups, role of, in the binding of nitroaromatic compounds to myeloma protein (immunoglobulin A) from mouse plasma-cell tumour MOPC 315 cells (Gettins, P., Givol, D. & Dwek, R. A.) 713–722

Nitroaromatic compounds, role of nitro groups in the binding of, to myeloma protein (immunoglobulin A) from mouse plasma-cell tumour MOPC 315 cells (Gettins, P., Givol, D. & Dwek, R. A.) 713–722

Nitrogenase complex, *Klebsiella pneumoniae*, electron-paramagnetic-resonance evidence for the formation of transient complexes of acetylene and ethylene with (Lowe, D. J., Eady, R. R. & Thorneley, R. N. F.) 277–290

Nitrogenase complexes, bacterial, electron-paramagnetic-resonance studies on the redox properties of the molybdenum–iron protein component of, between +50 and −450 mV (O'Donnell, M. J. & Smith, B. E.) 831–838

Nitrogenase system, *Spirillum lipoferum*, properties of (Ludden, P. W., Okon, Y. & Burris, R. H.) 1001–1003

Non-histone proteins, *see* Proteins, non-histone

Nuclei, L929-cell, mouse, involvement of deoxyribonucleic acid polymerase β in the biosynthesis of deoxyribonucleic acid in (Butt, T. R., Wood, W. M., McKay, E. L. & Adams, R. L. P.) 309–314

Nuclei, macro-, *Tetrahymena pyriformis*, composition of the histones of chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid from (Jones, R. W.) 155–164

Nuclei, macro-, *Tetrahymena pyriformis*, preparation of chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid from (Jones, R. W.) 145–153

Oligosaccharides, differences in the specificities of human liver β-*N*-acetylhexosaminidases detected by their activities towards, prepared from glycosaminoglycans (Bearpark, T. M. & Stirling, J. L.) 997–1000

Ovotransferrin, chicken-egg, derivation of equations relating the fractional saturation of *N*-terminal and *C*-terminal sites to the number of bound iron atoms per molecule of (Milne, J. G. C.) 541

Ovotransferrin, chicken-egg, iron-binding properties of (Williams, J., Evans, R. W. & Moreton, K.) 535–541

3-Oxo acid coenzyme A-transferase, kidney, sheep, purification and properties of (Sharp, J. A. & Edwards, M. R.) 759–765

Oxygen, mechanism of reaction of ferricyanide-pretreated mixed-valence-state ox heart mitochondrial membrane-bound cytochrome *c* oxidase with, at 173K (Clore, G. M. & Chance, E. M.) 811–820

Oxygen, mechanism of reaction of fully reduced ox heart mitochondrial membrane-bound cytochrome *c* oxidase with, at 176K (Clore, G. M. & Chance, E. M.) 799–810

Oxygen, reaction of, with *Pseudomonas aeruginosa* N.C.T.C. 6750 cytochrome *c*-551 oxidase (Greenwood, C., Barber, D., Parr, S. R., Antonini, E., Brunori, M. & Colosimo, A.) 11–17

Oxyhaemoglobin, human, mechanism of the breakdown of, on reaction with acetylphenylhydrazine (French, J. K., Winterbourn, C. C. & Carrell, R. W.) 19–26

Oxytocin, influence of length of the peptide chain on the formation of disulphide bonds in vasopressin and, and their analogues (Moore, G.) 403–409

Pancreas, ox, effects of method of preparation and storage on the allosteric behaviour of ribonuclease from (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 5–10

Pancreas, ox, nature of the allosteric interactions of ribonuclease from, with substrates and substrate analogues (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 1–4

Pancreas, ox, structure and enzyme activity of ribonuclease A from, after esterification at the glutamic acid-49 and aspartic acid-53 residues (Acharya, A. S., Manjula, B. N. & Vithayathil, P. J.) 821–830

Papain, use of 2,2′-dipyridyl disulphide in a spectrophotometric method for the detection of chymopapains as contaminants in preparations of (Baines, B. S. & Brocklehurst, K.) 345–347

Peptide, removal by transamination and scission of amino acid residues from, representing the cupric ion-transport site of human serum albumin (Sarkar, B., Dixon, H. B. F. & Webster, D.) 895–897

Peptides, biochemical applications of chemically induced nuclear polarization in phenols, catecholamines and (Muszkat, K. A., Weinstein, M. & Gilon, C.) 993–996

Periodate, oxidation by, and the configurations of glycosaminoglycans in solution (Scott, J. E. & Tigwell, M. J.) 103–114

Periphylla periphylla, *see* Jellyfish

pH, dependence on, of intrinsic fluorescence of chemically modified horse liver alcohol dehydrogenase (Parker, D. M., Hardman, M. J., Plapp, B. V., Holbrook, J. J. & Shore, J. D.) 269–275

pH, effects of ionic strength and, on the allosteric interactions of ox pancreatic ribonuclease with substrates and substrate analogues (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 1–4
Phenobarbital, substrate specificity and properties of liver uridine diphosphate glucuronyltransferase purified to apparent homogeneity from rats pretreated with (Burchell, B.) 749–757
Phenols, biochemical applications of chemically induced nuclear polarization in peptides, catecholamines and (Muszkat, K. A., Weinstein, M. & Gilon, C.) 993–996
Phenylglyoxal, inactivation by butane-2,3-dione, pyridoxal 5′-phosphate and, of *Bacillus megaterium* N.C.T.C. 10342 glutamate dehydrogenase and glutamate synthase (Hemmilä, I. A. & Mäntsälä, P. I.) 53–58
Pigeon, composition of the protein and lipid components of erythrocyte membrane from (Watts, C. & Wheeler, K. P.) 899–907
Pituitary gland, posterior, influence of length of the peptide chain on the formation of disulphide bonds in hormones biosynthesized by, and their analogues (Moore, G.) 403–409
Placenta, human, immunochemical characterization of human liver proteins structurally related to *N*-acetyl-β-D-glucosaminidase from (Carroll, M.) 191–196
Plasma-cell tumour MOPC 315 cells, mouse, role of nitro groups in the binding of nitroaromatic compounds to myeloma protein (immunoglobulin A) from (Gettins, P., Givol, D. & Dwek, R. A.) 713–722
Plasma, human, physical and chemical properties of α_2-macroglobulin from (Hall, P. K. & Roberts, R. C.) 27–38
Polydeoxyribonucleotides, bifunctional intercalation and sequence specificity in the binding of quinoxaline antibiotics to (Lee, J. S. & Waring, M. J.) 115–128
Polydeoxyribonucleotides, changes in mechanism and specificity related to structural changes in the interaction between synthetic analogues of quinoxaline antibiotics and (Lee, J. S. & Waring, M. J.) 129–144
Polypeptides, high-molecular-weight, spectrin, two, similarity of, from ox erythrocyte membrane (Dunn, M. J., Kemp, R. B. & Maddy, A. H.) 197–205
Polyuronides, oxidation by periodate and the configurations of, in solution (Scott, J. E. & Tigwell, M. J.) 103–114
Porphyra umbilicalis, determination of the amino acid sequence of ferredoxin from (Takruri, I., Haslett, B. G., Boulter, D., Andrew, P. W. & Rogers, L. J.) 459–466
Porphyrin carboxylic acids, use of high-pressure liquid chromatography with solvent systems containing phase-transfer agents for the direct determination of (Bonnett, R., Charalambides, A. A., Jones, K., Magnus, I. A. & Ridge, R. J.) 693–696
Procion Red HE-3B, immobilized, selective retardation of nicotinamide–adenine dinucleotide phosphate-dependent dehydrogenases on chromatography columns of (Watson, D. H., Harvey, M. J. & Dean, P. D. G.) 591–596
Proline, molecular weights and the problem of aggregation of enzymes involved in the biosynthesis of, in *Escherichia coli* (Hayzer, D. J. & Moses, V.) 219–228
Proline, potential errors arising from the use of a bioradiological procedure for assay of the biosynthesis of, by cell-free extracts of *Escherichia coli* (Hayzer, D. J. & Moses, V.) 207–217
Protease, *see* Proteinase
Protein, actin-like, stimulation by 1,25-dihydroxycholecalciferol of the incorporation of leucine into, in chick small-intestine brush border (Wilson, P. W. & Lawson, D. E. M.) 627–631
Protein, iron–sulphur, molybdenum-containing, *Desulfovibrio gigas*, determination of the redox potential of (Moura, J. J. G., Xavier, A. V., Cammack, R., Hall, D. O., Bruschi, M. & Le Gall, J.) 419–425
Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, skeletal-muscle, rabbit, evidence for an essential arginine residue in the recognition site of (Matsuo, M., Huang, C. & Huang, L. C.) 441–447
Protein, membrane (Band 3), erythrocyte, human, major, association of blood-group-I activity with (Childs, R. A., Feizi, T., Fukuda, M. & Hakomori, S.-I.) 333–336
Protein, molybdenum–iron, electron-paramagnetic-resonance studies on the redox properties of, of various bacterial nitrogenase complexes between +50 and −450 mV (O'Donnell, M. J. & Smith, B. E.) 831–838
Protein, myeloma (immunoglobulin A), role of nitro groups in the binding of nitroaromatic compounds to, from mouse plasma-cell tumour MOPC 315 cells (Gettins, P., Givol, D. & Dwek, R. A.) 713–722
Protein, retinoic acid-binding, skin, chick-embryo, purification and properties of (Sani, B. P. & Banerjee, C. K.) 643–649
Proteinase (actinidin), Chinese-gooseberry, determination of the amino acid sequences of peptides obtained by tryptic digestion of (Carne, A. & Moore, C. H.) 73–83
Proteinase, neutral, smooth-muscle, small-intestinal, rat, inactivation of native enzymes by (Beynon, R. J. & Kay, J.) 291–298
Proteinases, synovial-fibroblast and polymorphonuclear-leucocyte, rabbit, properties of, with activity towards proteoglycans (Werb, Z., Dingle, J. T., Reynolds, J. J. & Barrett, A. J.) 949–958
Protein–polysaccharides, *see* Proteoglycans
Proteins, composition of the lipids and, present in the pigeon erythrocyte membrane (Watts, C. & Wheeler, K. P.) 899–907
Proteins, different, synthesis of *N*-succinimidyl 3-(2-pyridyldithio)propionate and its use as a heterobifunctional reagent for the thiolation of proteins and for the conjugation of (Carlsson, J., Drevin, H. & Axén, R.) 723–737
Proteins, liver, human, immunochemical characterization of, structurally related to human placenta *N*-acetyl-β-D-glucosaminidase (Carroll, M.) 191–196
Proteins, myelin, sciatic-nerve, mouse, composition of, from Quaking and Jimpy mutants (Matthieu, J.-M.) 989–991
Proteins, non-histone, chromosomal, calf, lack of tissue specificity of (Rabbani, A., Goodwin, G. H. & Johns, E. W.) 497–505
Proteins, synthesis of *N*-succinimidyl 3-(2-pyridyldithio)propionate and its use as a heterobifunctional reagent for the thiolation of, and for the conjugation of different proteins (Carlsson, J., Drevin, H. & Axén, R.) 723–737

Proteins, wheat-kernel, monomeric and dimeric, structural studies of, with inhibitory activity towards yellow-mealworm α-amylase (Petrucci, T., Sannia, G., Parlamenti, R. & Silano, V.) 229–235
Proteoglycans, cartilage, laryngeal, pig, characterization by electric birefringence of, and of their complexes with hyaluronic acid (Isles, M., Foweraker, A. R., Jennings, B. R., Hardingham, T. & Muir, H.) 237–243
Proteoglycans, corneal, ox, different, interactions between (Speziale, P., Speziale, M. S., Galligani, L. & Balduini, C.) 935–939
Proteoglycans, properties of rabbit synovial-fibroblast and polymorphonuclear-leucocyte proteinases with activity towards (Werb, Z., Dingle, J. T., Reynolds, J. J. & Barrett, A. J.) 949–958
Pseudomonas aeruginosa N.C.T.C. 6750, oxidation by potassium ferricyanide of cytochrome *c*-551 oxidase from (Barber, D., Parr, S. R. & Greenwood, C.) 681–690
Pseudomonas aeruginosa N.C.T.C. 6750, reaction of oxygen with cytochrome *c*-551 oxidase from (Greenwood, C., Barber, D., Parr, S. R., Antonini, E., Brunori, M. & Colosimo, A.) 11–17
Pulmonary secretions, human, identification of hyaluronic acid in, of patients with asthma (Sahu, S. & Lynn, W. S.) 565–568
Pyridoxal 5′-phosphate, inactivation by butane-2,3-dione, phenylglyoxal and, of *Bacillus megaterium* N.C.T.C. 10342 glutamate dehydrogenase and glutamate synthase (Hemmilä, I. A. & Mäntsälä, P. I.) 53–58
1-Pyrroline-5-carboxylate reductase, *Escherichia coli*, molecular weights of, and other enzymes involved in the biosynthesis of proline and the problem of enzyme aggregation (Hayzer, D. J. & Moses, V.) 219–228
Pyruvate dehydrogenase complex, heart, pig, subunit stoicheiometry in the regulation by phosphorylation of the activity of (Sugden, P. H. & Randle, P. J.) 659–668
Pyruvate dehydrogenase complex, inactive, inhibition by alloxan-diabetes and starvation of the dephosphorylation of, to the active complex in rat heart mitochondria (Hutson, N. J., Kerbey, A. L., Randle, P. J. & Sugden, P. H.) 669–680

Quinomycin C, bifunctional intercalation and sequence specificity in the binding of, and other quinoxaline antibiotics to deoxyribonucleic acids (Lee, J. S. & Waring, M. J.) 115–128
Quinoxaline antibiotics, bifunctional intercalation and sequence specificity in the binding of, to deoxyribonucleic acids (Lee, J. S. & Waring, M. J.) 115–128
Quinoxaline antibiotics, changes in mechanism and specificity related to structural changes in the interaction between synthetic analogues of, and deoxyribonucleic acids (Lee, J. S. & Waring, M. J.) 129–144

Rainbow trout (*Salmo gairdneri*), isolation and partial characterization of serum lipoproteins and apolipoproteins of (Skinner, E. R. & Rogie, A.) 507–520
Red blood cells, *see* Erythrocytes
Redox potential, determination of, of the *Desulfovibrio gigas* molybdenum-containing iron–sulphur protein (Moura, J. J. G., Xavier, A. V., Cammack, R., Hall, D. O., Bruschi, M. & Le Gall, J.) 419–425
Reticulocytes, rabbit, retention at low temperatures of ribosomes from, on heparin bound to Sepharose 4B (Hradec, J. & Kříž, O.) 349–352
Reticulum, sarcoplasmic, skeletal-muscle, rabbit, determination of the amino acid sequences of cysteine-containing peptides obtained by tryptic digestion of calcium ion-transporting adenosine triphosphatase from (Allen, G. & Green, N. M.) 393–402
Retinoic acid-binding protein, skin, chick-embryo, purification and properties of (Sani, B. P. & Banerjee, C. K.) 643–649
Rhodospirillum rubrum N.C.I.B. 8255, kinetics of the reaction catalysed by ribulose bisphosphate carboxylase from (Christeller, J. T. & Laing, W. A.) 467–473
Ribonuclease A, pancreatic, ox, structure and enzyme activity of, after esterification at the glutamic acid-49 and aspartic acid-53 residues (Acharya, A. S., Manjula, B. N. & Vithayathil, P. J.) 821–830
Ribonuclease, pancreatic, ox, effects of method of preparation and storage on the allosteric behaviour of (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 5–10
Ribonuclease, pancreatic, ox, nature of the allosteric interactions of, with substrates and substrate analogues (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 1–4
Ribonucleic acid, ribosomal, composition of the histones of *Tetrahymena pyriformis* macronuclear chromatin containing deoxyribonucleic acid coding for (Jones, R. W.) 155–164
Ribonucleic acid, ribosomal, 5.8S, HeLa-cell, evidence on the conformation of, from the reaction of sodium bisulphite with specific cytidylate residues (Kelly, J. M., Goddard, J. P. & Maden, B. E. H.) 521–532
Ribonucleic acid, ribosomal, preparation of *Tetrahymena pyriformis* macronuclear chromatin containing deoxyribonucleic acid coding for (Jones, R. W.) 145–153
Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal
Ribosomes, reticulocyte, rabbit, retention at low temperatures of, on heparin bound to Sepharose 4B (Hradec, J. & Kříž, O.) 349–352
Ribulose bisphosphate carboxylase, *Rhodospirillum rubrum* N.C.I.B. 8255, kinetics of the reaction catalysed by (Christeller, J. T. & Laing, W. A.) 467–473

Saccharomyces cerevisiae, kinetics and mechanism of the reaction catalysed by potassium ion-activated aldehyde dehydrogenase from (Bostian, K. A. & Betts, G. F.) 787–798
Saccharomyces cerevisiae, rapid purification and properties of potassium ion-activated aldehyde dehydrogenase from (Bostian, K. A. & Betts, G. F.) 773–786
Salmo gairdneri, *see* Rainbow trout
Sarcoplasmic reticulum, *see* Reticulum, sarcoplasmic
Sciatic nerve, *see* Nerve, sciatic
Seminal fluid, human, use of affinity chromatography for the purification of lactate dehydrogenase isoenzyme X from (Kolk, A. H. J., Van Kuyk, L. & Boettcher, B.) 767–771
Serum albumin, *see* Albumin, serum
Serum lipoproteins, *see* Lipoproteins, serum
Serum transferrin, *see* Transferrin, serum
Sialoglycoproteins, anomalous gel-filtration behaviour of (Alhadeff, J. A.) 315–319

Skeletal muscle, *see* Muscle, skeletal
Skin, chick-embryo, purification and properties of retinoic acid-binding protein from (Sani, B. P. & Banerjee, C. K.) 643–649
Skin fibroblasts, human, cultured, secretion of isoenzymes of β-*N*-acetylglucosaminidase by (Willcox, P.) 433–439
Small intestine, *see* Intestine, small
Smooth muscle, *see* Muscle, smooth
Sodium dodecyl sulphate, solubilization by, of purified preparations of ox central-nervous-system myelin (Crang, A. J. & Rumsby, M. G.) 909–917
Sodium fluoride, solubilization of cow corpus luteum adenylate cyclase in Lubrol-PX, Triton X-100 or digitonin and the stabilizing effect of, present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924
Spectrin, membrane, erythrocyte, ox, similarity of the two high-molecular-weight polypeptides of (Dunn, M. J., Kemp, R. B. & Maddy, A. H.) 197–205
Spirillum lipoferum, properties of the nitrogenase system from (Ludden, P. W., Okon, Y. & Burris, R. H.) 1001–1003
Spleen, calf, lack of tissue specificity of the high-mobility group of chromosomal non-histone proteins from, and other tissues (Rabbani, A., Goodwin, G. H. & Johns, E. W.) 497–505
Spleen, human, properties of isoferritins from, and other tissues (Wagstaff, M., Worwood, M. & Jacobs, A.) 969–977
Starvation, inhibition by alloxan-diabetes and, of the dephosphorylation of inactive pyruvate dehydrogenase complex to the active complex in rat heart mitochondria (Hutson, N. J., Kerbey, A. L., Randle, P. J. & Sugden, P. H.) 669–680
Stomach, pig, comparison of the properties of intrinsic factor and cobalophilin present in digestive juices present in, and small intestine (Marcoullis, G., Merivuori, H. & Gräsbeck, R.) 705–712
Streptomyces clavuligerus, oxygenation of desacetoxycephalosporin C to desacetylcephalosporin C by 2-oxoglutarate-linked dioxygenases from *Acremonium chrysogenum* and (Turner, M. K., Farthing, J. E. & Brewer, S. J.) 839–850
Subcomponent C1q, complement, human, determination of the amino acid sequence of the *N*-terminal 108 amino acid residues of the B chain of (Reid, K. B. M. & Thompson, E. O. P.) 863–868
N-Succinimidyl-3-(2-pyridyldithio)propionate, synthesis of, and its use as a heterobifunctional reagent for the thiolation of proteins and for the conjugation of different proteins (Carlsson, J., Drevin, H. & Axén, R.) 723–737
Succinyl-coenzyme A–3-oxo acid coenzyme A-transferase, *see* 3-Oxo acid coenzyme A-transferase
Supernatant fraction, *see* Cytosol
Superoxide ion radicals, role of, in the breakdown of human oxyhaemoglobin on reaction with acetylphenylhydrazine (French, J. K., Winterbourn, C. C. & Carrell, R. W.) 19–26
Synovial fibroblasts, rabbit, properties of proteinases from, and polymorphonuclear leucocytes with activity towards proteoglycans (Werb, Z., Dingle, J. T., Reynolds, J. J. & Barrett, A. J.) 949–958
Tenebrio molitor, *see* Mealworm, yellow
Tetrahymena pyriformis, composition of the histones of chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid from macronuclei of (Jones, R. W.) 155–164
Tetrahymena pyriformis, preparation of chromatin containing deoxyribonucleic acid coding for ribosomal ribonucleic acid from macronuclei of (Jones, R. W.) 145–153
Thiolation, synthesis of *N*-succinimidyl-3-(2-pyridyldithio)propionate and its use as a heterobifunctional reagent for, of proteins and for the conjugation of different proteins (Carlsson, J., Drevin, H. & Axen, R.) 723–737
Thrombin, ox, effect of heparin on the inactivation of, by anti-thrombin III (Machovich, R. & Arányi, P.) 869–875
Thymus gland, calf, defined molecular-weight distribution of deoxyribonucleic acid from, after extensive sonication (Davis, A. W. & Phillips, D. R.) 179–183
Thymus gland, calf, general method for the purification of deoxycytidine kinase from, and various other mammalian tissues (Baxter, A., Currie, L. M. & Durham, J. P.) 1005–1008
Thymus gland, calf, lack of tissue specificity of the high-mobility group of chromosomal non-histone proteins from, and other tissues (Rabbani, A., Goodwin, G. H. & Johns, E. W.) 497–505
Thymus gland, chicken, rat and human, comparison of the compositions of glycosphingolipids and gangliosides of, and of chicken bursa of Fabricius (Narasimhan, R. & Murray, R. K.) 475–482
Toxin, cholera, time course of and rate-limiting steps in the activation by, of adenylate cyclase in rat liver (Fischer, J., Kohler, T. R., Lipson, L. G., Flores, J., Witkum, P. A. & Sharp, G. W. G.) 59–64
Transamination, removal by, and scission of amino acid residues from the peptide representing the cupric ion-transport site of human serum albumin (Sarkar, B., Dixon, H. B. F. & Webster, D.) 895–897
Transferrin, ovo-, chicken-egg, derivation of equations relating the fractional saturation of *N*-terminal and *C*-terminal sites to the number of bound iron atoms per molecule (Milne, J. G. C.) 541
Transferrin, ovo-, chicken-egg, iron-binding properties of (Williams, J., Evans, R. W. & Moreton, K.) 533–539
Transferrin, serum, human, binding of different iron donors to, and isolation of iron-binding fragments from the *N*-terminal and *C*-terminal regions of the protein (Evans, R. W. & Williams, J.) 543–552
Triostin A, bifunctional intercalation and sequence specificity in the binding of, and other quinoxaline antibiotics to deoxyribonucleic acids (Lee, J. S. & Waring, M. J.) 115–128
Triticum aestivum, *see* Wheat
Triton WR1339, effects of the administration of asialofetuin and, on the uptake by mouse liver of carcinoembryonic antigen and asialo-carcinoembryonic antigen from the blood (Thomas, P. & Jones, M.) 981–983

Triton X-100, solubilization of cow corpus luteum adenylate cyclase in Lubrol-PX, digitonin or, and the stabilizing effect of sodium fluoride present in the solubilization medium (Young, J. L. & Stansfield, D. A.) 919–924
Tropoelastin, aorta, chick-embryo, factors influencing the formation of isodesmosine and desmosine by the action of lysyl oxidase on (Narayanan, A. S., Page, R. C., Kuzan, F. & Cooper, C. G.) 857–862
Troponin C, skeletal-muscle, rabbit, characterization of a region of the primary amino acid sequence of, that is involved in calcium ion-dependent interaction with troponin I (Weeks, R. A. & Perry, S. V.) 449–457
Troponin I, skeletal-muscle, rabbit, characterization of a region of the primary amino acid sequence of rabbit skeletal-muscle troponin C that is involved in calcium ion-dependent interaction with (Weeks, R. A. & Perry, S. V.) 449–457

Uridine diphosphate glucose dehydrogenase, liver, ox, half-of-the-sites oxidation by 6,6′-dithiodinicotinate of (Franzen, J. S., Marchetti, P., Ishman, R., Ashcom, J. & Feingold, D. S.) 701–704
Uridine diphosphate glucuronyltransferase, liver, rat, substrate specificity and properties of, purified to apparent homogeneity from animals pretreated with phenobarbital (Burchell, B.) 749–757
Urine, human, use of high-pressure liquid chromatography with solvent systems containing phase-transfer agents for the direct determination of porphyrin carboxylic acids in, and other biological samples (Bonnett, R., Charalambides, A. A., Jones, K., Magnus, I. A. & Ridge, R. J.) 693–696

Vasopressin, influence of length of the peptide chain on the formation of disulphide bonds in oxytocin and, and their analogues (Moore, G.) 403–409
Vitamin B_{12}, comparison of the binding of, by intrinsic factor and cobalophilin present in digestive juices in different parts of the pig gastrointestinal tract (Marcoullis, G., Merivuori, H. & Gräsbeck, R.) 705–712
Vitamin D_3, *see* Cholecalciferol

Wheat (*Triticum aestivum*) kernel, structural studies of monomeric and dimeric proteins from, with inhibitory activity towards yellow-mealworm α-amylase (Petrucci, T., Sannia, G., Parlamenti, R. & Silano, V.) 229–235
Wool, sheep, determination of the amino acid sequences of peptides obtained by tryptic and chymotryptic digestion of a type-II α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of α-keratin from (Hogg, D. McC., Dowling, L. M. & Crewther, W. G.) 353–363
Wool, sheep, determination of the complete amino acid sequence of a type-I α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of α-keratin from (Gough, K. H., Inglis, A. S. & Crewther, W. G.) 373–385
Wool, sheep, determination of the complete amino acid sequence of a type-II α-helical segment of *S*-carboxymethylkerateine-A from the low-sulphur fraction of α-keratin from (Crewther, W. G., Inglis, A. S. & McKern, N. M.) 365–371
Wool, sheep, statistical analysis of the amino acid sequences of α-helical segments of *S*-carboxymethylkerateine-A from the low-sulphur fraction of α-keratin from (Elleman, T. C., Crewther, W. G. & Van der Touw, J.) 387–391

Yeast, baker's (*Saccharomyces cerevisiae*), kinetics and mechanism of the reaction catalysed by potassium ion-activated aldehyde dehydrogenase from (Bostian, K. A. & Betts, G. F.) 787–798
Yeast, baker's (*Saccharomyces cerevisiae*), rapid purification and properties of potassium ion-activated aldehyde dehydrogenase from (Bostian, K. A. & Betts, G. F.) 773–786

Zinc ions, role of, in the stabilization of the dimeric form of *Bacillus amyloliquefaciens* α-amylase (Tellam, R., Winzor, D. J. & Nichol, L. W.) 185–190